



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Analysis of the Dynamics, Outcome, and Prerequisites of the first German SARS-CoV-2 Superspreading Event

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-059809
Article Type:	Original research
Date Submitted by the Author:	07-Dec-2021
Complete List of Authors:	Wessendorf, Lukas; University Hospital Bonn, Virology Richter, Enrico; University Hospital Bonn, Virology Schulte, Bianca; University Hospital Bonn, Institute for Virology Schmithausen, Ricarda Maria; University Hospital Bonn, Hygiene Exner, Martin; University Hospital Bonn Lehmann, Nils; University of Duisburg-Essen, Institute for Medical Informatics, Biometry and Epidemiology Coenen, Martin; Rheinische Friedrich Wilhelms Universität Bonn, Clinical Study Core Unit, Study Center Bonn (SZB) Fuhrmann, Christine; University Hospital Bonn Kellings, Angelika; University Hospital Bonn Hüsing, Anika; University Hospital Essen Jöckel, Karl-Heinz; Institute of Medical Informatics, Biometry and Epidemiology, University Duisburg- Essen Streeck, Hendrik; German Center for Infection Research
Keywords:	COVID-19, Infection control < INFECTIOUS DISEASES, VIROLOGY

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Analysis of the Dynamics, Outcome, and Prerequisites of the first German SARS-CoV-2 Superspreading Event

Lukas Wessendorf¹, Enrico Richter¹, Bianca Schulte¹, Ricarda Maria Schmithausen², Martin Exner², Nils Lehmann³, Martin Coenen⁴, Christine Fuhrmann⁴, Angelika Kellings⁴, Anika Hüsing³, Karl-Heinz Jöckel³, and Hendrik Streeck^{1,*}

¹ Institute of Virology, University Hospital, University of Bonn, Germany, and German Center for Infection Research (DZIF), partner site Bonn-Cologne

² Institute of Hygiene and Public Health, University Hospital, University of Bonn, Germany

³ Institute of Medical Informatics, Biometry und Epidemiology (IMIBE), University Hospital Essen, Germany

⁴ Clinical Study Core Unit, Study Center Bonn (SZB), Institute of Clinical Chemistry and Clinical Pharmacology, University Hospital Bonn, Germany

* Corresponding author: Hendrik Streeck, Institute of Virology, University Hospital Bonn, Venusberg-Campus 1, 53127 Bonn, Germany, Hendrik.Streeck@ukbonn.de

Keywords: SARS-CoV-2, COVID-19, superspreading, cohort study, ventilation

Word count: 3903

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

49 **Abstract**

50

51 **Objectives**

52 Determining the SARS-CoV-2 RT-PCR positivity rate, SARS-CoV-2-specific antibody

53 levels of the participants, and analyzing the conditions and dynamics of

54 superspreading, including ventilation, setting dimensions, distance from infected

55 persons and behavioral patterns.

56

57 **Design**

58 51 days after the event all participants were asked to give blood samples, pharyngeal

59 swabs and answer a self-administered questionnaire. Metric room coordinates for all

60 tables, seats, and ventilation-points were assessed.

61

62 **Setting**

63 The superspreading event took place during festivities including 450 people (6-79

64 years of age) in a building of 27 m x 13.20 m x 4.20 m.

65

66 **Participants**

67 All persons who took part in the event which led to superspreading of SARS-CoV-2

68 were invited to participate in this study.

69

70 **Interventions**

71 No interventions were performed.

72

73 **Primary and Secondary Outcome Measures**

74 The primary outcome measure was infection status by combining RT-PCR results with

75 ELISA results. Secondary outcomes were symptoms as stated in the questionnaire.

76

77 **Conclusions**

78 We analyzed infection rates and risk of infection depending on age, alcohol

79 consumption, and ventilation. Overall, 46% of participants had been infected. Spatial

80 distribution of infected participants was associated with proximity to the ventilation

81 system (OR 1.39, 95% KI [0.86; 2.25]). The risk of infection was highly associated with

82 age, thus children (OR: 0.33 [0.267; 0.414]) and young adults had a lower risk than

older participants resulting in an average infection risk increase of 28% per 10 years age difference. Behavioral differences reduced the risk of infection including time spent outside (OR: 0.55 [0.33; 0.91]) or smoking (OR: 0.32 [0.124; 0.81]).

Strengths and Limitations of this Study

- Strength: The setting and the participant group are extremely well-defined.
- Weakness: Some participants left the venue during the event for short times.
- Strength: Participants were invited by only one criteria, namely their presence at the superspreading event; no other preselection/bias took place during enrollment.
- Weakness: The event size was below 1000 people (450), therefore it was not possible to recruit more than 411 study participants.

Article Summary

The scientific literature was searched for the term "superspreading event AND Covid-19 OR Sars Cov 2" and identified published papers from China, South Korea, Europe, and North America. Most researchers analyzed superspreading events within a health care setting e.g. in hospitals or nursing homes, or described the general impact of superspreading events on the global pandemic. Only a few metanalyses of transmission clusters analyzed party occasions (e.g. a nightclub in Berlin, Germany) as superspreading events. These reports describe less than 100 infections and are very limited due to missing data or reporting biases. Therefore, the ability to draw scientific conclusions is also limited. Additionally, to our knowledge, there are no studies, which investigated individual behavior, the location, and role of children during a superspreading event. The research for the study started April 2020 and was concluded in June 2021.

Our report analyzes the first COVID-19 superspreading event in Germany in detail, which was not only a unique setting but also included children and adults in the same room. We demonstrate that nearly half of the participants were infected with SARS-CoV-2 and that the proximity of the seating to the ventilation system was an important risk factor for infection. The data showed that low physical distance including singing and duration of attendance at this event increased the risk of infection, while regular

smoking and spending the break of the event outside lowered the risk of infection. This underlines the benefit of airing to lower the amount of both droplets and aerosols. Furthermore, we found lower infection in children than adults despite being in the same room suggesting differences in infectability in children. Indeed, we observed that an additional 10 years of age is on average associated with 28% increased risk of infection.

Taken together, the results demonstrate the importance of the ventilation system during superspreading events. In particular children and young adults had a lower risk of infection during the event indicating that they have a limited role during this pandemic. Overall, our data demonstrate in detail age-dependent infectability as well as highlights to understand transmission dynamics in order to improve comprehensive public health preparedness measures.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a highly transmissible and pathogenic RNA virus that emerged in late 2019 and has caused a pandemic threatening human health and public safety worldwide.¹ While factors shaping the dynamics of a pandemic are multifactorial, virulence and reproductive number are important properties of a virus.² For SARS-CoV-2 there is a substantial over-dispersion of the secondary infection distribution (individual R_0) for an individual infected with SARS-CoV-2.² An over-dispersed R_0 means that most infected people do not transmit (individual $R_0 = 0$) while a minority of infected people are super-spreaders (individual $R_0 > 5$). Superspreading has been observed for many infectious pathogens, such as measles or SARS.³ During the SARS pandemic in 2003 a superspreading event was defined as one infected person infecting eight others.⁴ For SARS-CoV-2 it has been estimated that 80% of the infections are caused by 10% of infected individuals highlighting the importance of the cluster factor (k).² In Germany an indoor carnival event in the beginning of 2020 is considered as the first major outbreak in a German city and was considered a hotspot during the beginning of the pandemic in Germany.⁵ Other SARS-CoV-2 superspreading events worldwide have been linked to indoor gatherings with close proximity of individuals.⁶ Nevertheless,

most of the reported superspreading events had less than 100 cases and the reports are limited by missing data or a reporting bias.⁶

Here, we closely examined the prerequisite of a unique super-spreading event in Germany during the SARS-CoV-2 pandemic, where nearly half of the participants became infected including children. We systematically analyzed infection rate, potential individual, and environmental risk factors for infection as well as the role of the ventilation system.

Materials and methods

Study design and sampling

This cross-sectional epidemiological study was conducted 51 days after a carnival celebration in the beginning 2020. Eleven days after the event authorities sent all known participants into quarantine after testing 38 out of 99 individuals PCR-positive. All adults known to have attended the event were invited to participate in the study. About 450 persons attended the event of which 411 participated in the study (**figure 1**, participation rate 91.3%). All study participants provided written informed consent before enrolment.

Self-administered questionnaires included questions about demographic background, symptoms of viral infection as well as detailed information about the behavior during the event. Participants' arrival and exit times were assessed in 1-hour categories. Study participants were asked to provide blood specimens and pharyngeal swabs for further analysis.

Research Ethics Approval

The Ethics Committee of the Medical Faculty of the University of Bonn approved the study (approval number 085/20).

Patient and Public Involvement Statement

The concept and organization of the study on-site was in close cooperation and agreement with the local administration, the county commissioner and the society-leader of the carnival event. Additionally, the community was in need of SARS-CoV-2

testing, because at this time the availability of testing was still limited, therefore they invited the collaboration with our team of scientists and physicians.

Pharyngeal swab and blood preparation

Pharyngeal swabs of participants were performed with FLOQSwabs (Copan) and immediately stored in UTM RT-mini tubes containing UTM Viral Stabilization Media (Copan) at 4 °C. Venous blood was drawn into EDTA tubes (Sarstedt) per volunteer and was transported to the laboratory at the University Hospital Bonn.

Anti-SARS-CoV-2 ELISA

Anti-SARS-CoV-2 IgA and IgG were determined using enzyme-linked immunosorbent assays (ELISA) on the EUROIMMUN Analyzer I platform.⁵ According to the manufacturer's instructions a result was considered positive when a ratio (extinction of sample/extinction of calibrator) of 0.8 or higher was reached. The guidelines of the German Medical Association (RiliBÄK) were abided by, including internal and external quality controls.

Reverse transcription polymerase chain reaction (RT-PCR)

Viral RNA was extracted from each 300µl swab sample via the chemagic Viral 300 assay (according to manufacturer's instructions) on the Perkin Elmer chemagic™ Prime™ instrument platform. The presence of two viral target genes (E and RdRP) was assessed in each sample by real time RT-PCR (SuperScript™III One-Step RT-PCR System with Platinum™ TaqDNA Polymerase, Thermo Fisher). The following primers were used, for E gene: E_Sarbeco_F1 and R, and probe E_Sarbeco_P1, for RdRP gene: RdRP_SARSr_F, and R, and probe RdRP_SARSr-P2.⁷ In addition, an internal control for RNA extraction, reverse transcription, and amplification was applied to each sample (innuDETECT Internal Control RNA Assay, Analytik Jena #845-ID-0007100). If amplification occurred in both virus-specific reactions samples were considered positive.

SARS-CoV-2 Neutralization Assay

A plaque reduction neutralization test was used to determine SARS-CoV-2 neutralization capacity as previously described.⁵ Briefly, plasma samples were heat-

inactivated and supernatant transferred to a new tube and serially two-fold diluted in OptiPROTMSFM (Gibco) performed. 120 mL of each plasma dilution was mixed with 80 plaque-forming units (PFU) of SARS-CoV-2 in 120 mL OptiPRO SFM (GIBCO) cell culture medium and seeded with 1.25×10^5 Vero E6 cells/well. Subsequently, the inoculum was removed and cells were overlaid with a mixture of carboxymethylcellulose (Sigma) and 2xMEM (Biochrom). Following 3-day incubation, the overlay was removed and the 24 well plates were fixed using a 6% formaldehyde solution and stained with 1% crystal violet in 20% ethanol revealing the formation of plaques. Finally, the neutralizing titers were calculated as the reciprocal of serum dilutions resulting in neutralization of 50% input virus (NT50), read out as reduction in the number of plaques.

Data management and quality control

The Clinical Study Core Unit of the Study Center Bonn (SZB) supported the study by outlining the study protocol and developing the informed consent form as well as participants information sheets with respect to data management and quality control. The data were gathered on paper-based Case Report Forms (pCRF). Data was entered as double-data-entry into the REDCap study database programmed and hosted by SZB. Study personnel was trained by experienced members of the SZB. A quality manager was on site to support the study team. Monitoring of trial data and informed consent forms was performed according to the monitoring plan by qualified SZB staff. The ethics committee of the Medical Faculty of the University of Bonn was involved and approved the study (reference no. 085/20)

Spatial information

Metric room coordinates (length and width [m]) for areas, tables, seats and ventilation shafts were assessed via measurements, seating plan and photos from the event. Persons providing multiple positions were considered as spending an equal amount of time on different positions. When exact seating was unclear and information was available on table or greater area localisation (bar, stage), average coordinate values were used.

On the grounds of these coordinates, we calculated pairwise metric distances between all persons and distances to closest inletting and purging airshafts. For all persons their pairwise inverse distances were summarized as mean inverse distance. Inverse metric

distances to persons or airshafts were regarded as representing infectious potential through local proximity, and inverse distances were capped at 2.5 (the inverse of the width of a seat of 0.4 m). Alternatively, we counted all, and all infected persons within adjacent rings of 1.5 m width around each participant as a measure of crowdedness and infectious potential.

Statistical analysis

Associations between positive infection status and exposure variables were analysed via logistic regression models. Exposure variables were included crudely, and adjusted for potential confounding factors age, sex, and duration of attendance as fixed effects. To correct for common household effects a random effects model was used. We present odds ratios with 95% confidence intervals. Because we present data on a single specific event among a limited number of participants, we completely refrain from presenting p-values. All analyses were done with SAS 9.4.

Results

411 out of estimated 450 participants of the event responded to our study invitation, resulting in a response rate of 91.3%. 404 individuals provided plasma samples and 316 pharyngeal swabs (figure 1). Genders were represented equally among all 404 participants (48% were male) with a broad range in age ((range 6-79) median age 36 years) and level of education (table 1). 297 individuals were residents of the community the event took place in, 103 lived in other parts of the county, and 11 were external visitors.

Overall, 186 out of 404 individuals tested seropositive for IgG- and 161 for IgA-antibodies (suppl. figure 1). To confirm seropositivity we performed a plaque reduction neutralization assay (suppl. figure 2) demonstrating neutralizing activity against SARS-CoV-2 of their respective antibody responses. Given the low specificity of the IgA assay, IgA seropositivity was not further considered. As we tested for seropositivity 51 days after the superspreading event, we additionally performed SARS-CoV-2 RT PCR analysis from pharyngeal swabs to exclude potential recent infections. Indeed, 19 participants tested positive in RT-PCR, and were therefore not considered in the study as there was no likelihood of infection during the superspreading event.

	Not infected N%	Infected N%
Total number	218	186
Female	114 (52%)	89 (48%)
Age		
<18 years	31 (14%)	15 (8%)
18-24 years	30 (14%)	20 (11%)
25-39 years	81 (37%)	43 (23%)
40-64 years	71 (33%)	100 (54%)
65+ years	5 (2%)	8 (4%)
BMI (kg/m²) (std dev)	24.3 (5.12)	26.2 (5.16)
Participating house hold member (std dev)	2.1 (1.12)	2.4 (1.16)
Highest level of formal education		
None	27 (13%)	13 (7%)
Lower secondary school	27 (13%)	23 (13%)
Secondary school	55 (26%)	71 (39%)
Higher education entrance qualification	54 (25%)	34 (18%)
(Technical) university degree	52 (24%)	43 (23%)
Duration of attendance [h] (std dev)	4.7 (2.06)	5.8 (1.85)
Service team	4 (2%)	22 (12%)
On stage during the event	80 (37%)	62 (34%)
Member of the 'Council of 11'	6 (3%)	18 (10%)
On stage during 'finale'	26 (12%)	48 (26%)
Behavior during break		
Remaining seated	73 (36%)	85 (48%)
Going outside	114 (55%)	72 (39%)
Alcohol consumption [drink] (std dev)	11.3 (7.76%)	12.2 (7.40)
Former smoker	34 (16%)	45 (24%)
Active smoker (>10 cigarettes per day)	54 (25%)	23 (12%)
At least one comorbidity	29 (13%)	28 (15%)
Avg. distance to other participants [m] (std dev)	9.2 (1.68)	9.1 (1.70)
Distance to air inlet [m] (std dev)	6.1 (3.22)	6.0 (3.30)
Distance to air outlet [m] (std dev)	4.8 (2.94)	5.1 (2.87)

Table 1: Distribution of demographic factors and exposure information of interest among study participants who tested positive or negative in serology test of SARS-Cov2-infection.

1
2
3 285 Overall, we found that 46.0% (95% CI: [41.2%; 51.0%]) tested seropositive who
4
5 286 attended the event, which was significantly higher than the overall estimated infection
6
7 287 rate in the same community at large at that time. Indeed, officially 3.1% of the
8
9 288 community were reported as positive cases at that time and we estimated the infection
10 289 rate as 15.5% (95% CI:[12.3%; 19.00%])**Error! Bookmark not defined.** for the
11
12 290 community. Taken together, an estimated 46% of participants became infected during
13
14 291 a single superspreading event.

15 292
16 293 No association between sex and risk of infection was found ((OR: 1.01 [0.65; 1.58]) for
17
18 294 women). On average infected individuals had a higher body mass index (26.2kg/m²
19
20 295 compared to 24.3kg/m² for uninfected individuals). Infected participants were more
21
22 296 likely to be clustered living in the same household (**table 1**). Having at least one
23
24 297 comorbidity, including lung disease (42.3%), cardiovascular disease (53.3%),
25 298 neurological disease (16.7%), cancer (58.3%) or diabetes (80%), did not increase the
26
27 299 risk of infection (OR: 0.64 [0.33; 1.26]). In conclusion, sex and comorbidities did not
28
29 300 seem to affect the risk of infection. We next assessed whether age influenced the risk
30
31 301 of infection at the event, considering sex, duration of attendance and common
32
33 302 household as covariates. Comparison across age-categories showed a lower risk for
34
35 303 children (OR: 0.31 [0.14; 0.69]), and also for young adults (18-25 years, OR: 0.53 [0.26;
36
37 304 1.09]) as well as adults between 25 and 40 years (OR: 0.48 [0.28; 0.85]) in comparison
38
39 305 to older adults (40 to 65 years) (OR: 1, reference), while seniors had a slightly higher
40
41 306 risk (older than 65 years, OR: 1.1 [0.31; 3.97]) (**figure 3**). Our data suggest that an
42
43 307 additional 10 years of age are on average associated with 28% increased risk of
44
45 308 infection (OR: 1.28 [1.10; 1.48]).

46 309
47 310 To understand the spreading dynamics of SARS-CoV-2 during the event, we first
48
49 311 performed a detailed analysis of potential risk factors and social behavior. The event
50
51 312 consisted of speeches, dance, and music performances for a total of five hours, with
52
53 313 one large intermission and was hosted at a small community center (320 square
54
55 314 meters) with a stage up front and a bar in the back close to the entrance. Alcoholic and
56
57 315 nonalcoholic drinks were served in glasses and a food truck was located outside in
58
59 316 front of the venue. While most participants were sitting in the hall, a committee of
60
317 eleven individuals hosting the event were sitting on stage. The eleven people on stage
318 switched after a break. With approximately 450 participants there were about 1.4
319 individuals per square meter and the tables, each with two benches, were arranged in

two blocks with an alley to the stage (**figure 2**). Infected participants had been seated mostly at tables close to the bar, at the bar, or on stage. One table with 8 out of 11 infected people, was located far away from the bar at the other side of the hall and close to an air inlet. The group sitting on stage showed high numbers of infection (18 infected out of 24, **table 1**). We first analyzed whether the ventilation system influenced the distribution of SARS-CoV-2 infected individuals. It is important to state that the system's air flow consisted of 75% used and 25% fresh air. The air flow can be described as clockwise. The air system uses vents along one side of the venue and on stage to take in air (**figure 2**, air inlets purple). After 25% of fresh air has been added and the air has been filtered, vents along the other side of the venue return the air into the room (**figure 2**, air outlets blue). All ventilation points received the same amount of air due to throttle valves. For noise protection reasons windows remained closed. The air-system used F7-Filters ($\text{ISO ePM} \geq 2,5$) and had an air volume flow of 7500 m³/h.

Most tables located close to the air-inlets and showed no or only few infections (**figure 2**, green) also most surrounding tables showed low numbers of infection (**figure 2**, yellow). Tables close to the air-outlets (**figure 2**) show high (4 or 5 infected per table) and very high (6 or 7 infected per table) numbers of infected individuals. It is important to mention that the overall number of participants per table was not equal for all tables. Greater proximity to air outlets was associated with increased risk of infection with a crude OR=1.39 [0.86; 2.25]. This association remained stable and was hardly attenuated from adjustment for proximity to air inlet, age, gender, duration of attendance, proximity to other infected persons, stage-activity and going outside during the intermission (**figure 4**, multiple adjusted OR=1.26 [0.63; 2.50]). A similar apparent effect for proximity to air inlets (crude OR=1.17 [0.72; 1.89]) disappeared when duration of attendance was added to the model (**figure 4**, multiple adjusted OR=1.01 [0.53; 1.94]). Overall, however, we found the increased risk for individuals located closer to the air outlet remarkably persistent (**figure 4**).

We further studied the sum of the inverse distance to all infected participants as a measure of proximity to either one common virus source or mutual infection. However, there was no evidence for increased risk of infection from greater proximity to other infected persons (**suppl. table 1**). Furthermore, we found no evidence for a single

person being the source of the infection using 401 quantile-plot analysis conducted for each participant as potential source of infection separately (**suppl. figure 3**).

To understand the association of risk with behavior patterns we next investigated the influence of several factors on SARS-CoV-2 infection including time spent outside, smoking, performing on stage and participation during the final act (“Finale”) for 30 minutes. Results were all adjusted for age, sex, common household, and duration of attendance. Participation in multiple performances did not increase the risk of infection (OR per performance: 1.08 [0.91; 1.27]) while participation in the last “Finale” indicated a trend towards increased risk of infection (OR: 1.41 [0.65; 3.02]) (**figure 4**). Duration of attendance was persistently and strongly associated with an increased infection risk of 32% with each additional hour spent at the party (OR per hour: 1.32 [1.16; 1.49]). All further analyses were adjusted for this variable as potential confounding factor.

We next determined the level of alcohol consumption as number of drinks (high-proof liquor or beer) and did not observe any influence for the amount of alcohol consumption on the risk of becoming infected (OR per drink: 1.00 [0.96; 1.05]). Furthermore, participants who spent the break outside were less likely to be infected (OR: 0.55 [0.33; 0.91]) compared to individuals who spent the break inside the venue hall (**figure 4**). Interestingly, however, when we determined the impact of being regular smoker (defined as smoking of at least 10 cigarettes a day) on the risk of SARS-CoV-2 infection we observed a reduced risk of infection (OR: 0.32 [0.12; 0.81]) even after adjustment for “time spent outside”. In conclusion, duration of attendance at the carnival party increased the risk of infection, the number of alcoholic drinks was not associated with infection risk, while regular smoking and spending the break of the event outside lowers the risk of infection.

We next stratified seropositive individuals by their reported symptoms. Odds-ratios for each symptom were calculated for the timespan of 14 days following the event (**figure 5**). Similar to previous reports⁸ loss of smell (OR: 8.78 [4.81; 16.02]) and taste (OR: 10.09 [5.13; 19.88]) were strongest associated with SARS-CoV-2 infection. Other symptoms which were strongly associated with COVID-19 were: sweats and chills (OR: 5.28 [3.08; 9.07]), muscle and joint ache (OR: 5.19 [3.19; 8.44]), fatigue (OR: 4.22 [2.76; 6.45]) and fever (OR: 3.73 [2.10; 6.63]) (**figure 5**). Importantly, 15.1% of

the infected individuals reported no symptoms at all in a period of 14 days after the event. The rate of asymptomatic infections of participants of the event was lower than generally observed in the community the event took place in (36%).⁵ Overall, there was a lower proportion of asymptomatic cases among individuals infected after the event compared to members of the community, while loss of smell and taste showed the strongest association with an infection.

Discussion

The high overdispersion characteristics of SARS-CoV2 and its ability to be transmitted via aerosols under certain conditions are one of the main reasons that the beginning of the SARS-CoV2 pandemic was shaped by superspreading events.^{9,10} Germany's first superspreading event was an indoor carnival event in the beginning of 2020 in a rural the community. In this naturally occurring experiment, we demonstrate that nearly half of the participants became infected and demonstrate multiple prerequisites of such an event and risk factors for becoming infected. While our study population is not a representative sample of the general population the event may be regarded as exemplary for similar party occasions and may help reduce the number of infected in the future.

An important factor associated with infection risk was the ventilation system and the individual proximity to the ventilation outlets. Individuals close to the air-outlets that contained air with low amount of fresh air had the highest infection risk compared to those close to the air-inlets. This is in line with previous studies that demonstrated SARS-CoV-2 to be able to become air-borne under certain conditions and that the ventilation system can have an influence on virus spread.^{11,12,13} The air filters in the venue were not capable of intercepting virus particles supporting the notion on the importance of proper indoor ventilation systems.^{14,15} Indeed, spending the break of the event outside decreased the possibility of infection underscoring the benefit of proper ventilation to lower the amount of aerosols. Due to the nature of the event, the spatial distribution of the participants was not fixed throughout the evening, and not perfectly recapitulated, so this information carries some error. However, allowing for multiple positions per person we used all available information. Assuming further error in the spatial data to be random, this might lead to a dilution of effects, i.e. true associations

1
2
3 421 may remain undetected. Complementary analyses including e.g. the persons'
4 422 functions during the event show consistent results, so we see no evidence suggesting
5 423 bias in our findings.

6 424
7
8 425 The consumption of alcoholic drinks did not increase the risk of infection. While it has
9 426 been assumed that the alcoholic effect of decreased social inhibition may increase
10 427 likelihood of infection, we did not find any evidence for this association questioning
11 428 measures of a ban on alcohol to reduce numbers of infected. It is known that current
12 429 and former smokers disproportionately suffer from severe COVID-19 and their
13 430 numbers are relatively increased among those patients that need intensive care
14 431 treatment compared to non-smokers.^{16,17} However, it has been previously speculated
15 432 that the risk of infection is lower for smokers.¹⁸ Furthermore, a meta-analysis of seven
16 433 studies suggests that smokers have a reduced risk of testing positive for SARS-CoV-
17 434 2.¹⁹ Interestingly, we also observed a protective effect for an infection with SARS-CoV-
18 435 2, thus our findings support those statements and show an even greater protective
19 436 effect. The association might for example be explained by a role of the nicotinic
20 437 acetylcholine receptor.²⁰ While we strongly advise that smoking should not be
21 438 considered as a protective habit to prevent risk of infection, this knowledge may lead
22 439 to the investigation of a therapeutic or prophylactic treatment on the basis of this
23 440 molecular target.²¹

24 441
25 442 Our results indicate a trend that younger people are less likely to be infected compared
26 443 to older age groups. This trend is strongest for people under 18 but levels out over 40
27 444 years of age. The risk of infection for children in superspreading events has not been
28 445 investigated but the overall risk for infection in children seems to be lower than for
29 446 adults as a systematic review and its recent update reported, which is further supported
30 447 by our findings.^{22,23} As all individuals were exposed at the same event and time our
31 448 study is a perfect model for the previously described notion, that children are less likely
32 449 to become infected. Indeed, a recently published meta-analysis by Viner et al. showed
33 450 a low susceptibility for children and adolescents (OR of 0.56 (95% CI, 0.37-0.85))
34 451 which strongly supports our findings of a lower risk of infection in that age group, which
35 452 is even lower in our study.²⁴ Our finding supports the previously shown subordinate
36 453 influence on the spreading of the virus by children. The finding that each 10 years of
37 454 age increase the risk of infection during an event indicates that younger people and

their limited role should be considered when measures to contain the pandemic are implemented. Taken together, we could demonstrate important risk factors for infection during a superspreading event, which helps to understand transmission dynamics in order to improve comprehensive public health preparedness measures.

Acknowledgments

We would like to thank the inhabitants of the town in question for their participation. We would also like to thank the local government of the town where the outbreak took place for their support to conduct the study. Furthermore, we would like to thank the following people who helped with the study: Rudi Kirschen, Gero Wilbring, Janett Wieseler, Marek Korencak, Ryan Natrass, Jernej Pusnik, Maximilian Becker, Ann-Sophie Boucher, Marc Alexander de Boer, Rebekka Dix, Sara Dohmen, Kim Friele, Benedikt Gansen, Jannik Geier, Marie Gronemeyer, Sarah Hundertmark, Nora Jansen, Michael Jost, Louisa Khorsandian, Simon Krzycki, Ekaterina Kuskova, Judith Langen, Silvia Letmathe, Ann-Kathrin Lippe, Jonathan Meinke, Freya Merker, Annika Modemann, Janine Petras, Sophie Marie Porath, Anna Quast, Laurine Reese, Isabel Maria Rehbach, Jonas Richter, Thea Rödiger, Eva Schmitz, Tobias Schremmer, Louisa Sommer, Jennifer Speda, Yuhe Tang, Oliver Thanscheidt, Franz Thiele, Johanna Thiele, Julia Tholen, Moritz Transier, Maïke van der Hoek, Tillmann Verbeek, Sophia Verspohl, Kira Vordermark, Julian Wirtz, Marina Wirtz, Lisa Zimmer, Philip Koenemann, Adi Yaser, Lisa Anna, Katharina Bartenschlager, Lisa Baum, Roxana Böhmer Romero, Diana de Braganca, Isabelle Engels, Moritz Färber, Carina Fernandez Gonzalez, Lucia Maria Goßner, Victoria Handschuch, Franziska Georgia Liermann, Steffan Meißner, Laura Racenski, Patrick, Denis Raguse, Larissa Reiß, Maximilian Rölle, Franziska Scheele, Chiara Schwippert, Arlene Christin Schwippert, Antonia Seifert, Joshua David Stockhausen, Sofia Waldorf, Leonie Weinhold, Nicolai Trimpop, Julia Reinhardt, Vera Gast, Michelle Yong, Eva Engels, Jonathan Meinke, Susanne Schmidt, Janine Schulte, Saskia Schmitz, Kübra Bayrak, Regina Frizler, Katarzyna Andryka, Sofía Soler, Bastian Putschli, Thomas Zillinger, Marcel Renn, Patrick Müller, Dillon Corvino, Zeinab Abdullah, Katrin Paeschke, Hiroki Kato, Florian Schmidt, Maximilian Baum, Celina Schüter, Daniel Hinze, Martina Schmidt, Arcangelo Ricchiuto, Sonja Gross, Uta Wolber, Tobias Höller, Marion Zerlett, Esther Sib, Benjamin Marx, Souhaib Aldabbagh. We thank the Medizinischer Dienst der

Krankenkasse (MDK) Nordrhein for their generous help in conducting this study. In particular we would like to thank: Tanja Bell, Ina de Hesselle-Taddey, Susanne Goll, Katja Hoffmann-Pruss, Niko Kalamakidis, Verena Kayser, Hildegard Kessler, Norbert Körfer, Daniela Kroll, Florian Messerschmidt, Svenja Peters, Sebastian Schröder, Linda Schroers, Michael Zimmermann, Klaus-Peter Thiele. We thank Stefan Holdenrieder, Alexander Semaan, Bernd Pötzsch, and Georg Nickenig for providing control samples. The idea, the plan, the concept, protocol, the conduct, the data analysis, and the writing of the manuscript of this study were independent of any third parties, including the government of North Rhine-Westphalia, Germany.

Author Contributions

L.W. wrote the manuscript. L.W., RMS, and ER organized and ran the testing center. ER and BS organized and performed sample processing, analysis, and corrected the manuscript. NL and AH performed statistical analysis. MC, CF, and AK monitored the study. ME, KHJ and H.S oversaw the study and corrected the manuscript.

Data statement

All data from the study will be made available upon reasonable request.

Funding

The government of North Rhine-Westphalia (Germany) supported the study with 65,000 Euro.

Conflicts of interest

The authors declare no conflict of interest.

References

1. Chakraborty I, Maity P. COVID-19 outbreak: Migration, effects on society, global environment and prevention. *Sci Total Environ* 2020; **728**: 138882.

2. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. *Nature* 2005; **438**(7066): 355-9.

3. De Serres G, Markowski F, Toth E, et al. Largest measles epidemic in North America in a decade--Quebec, Canada, 2011: contribution of susceptibility, serendipity, and superspreading events. *J Infect Dis* 2013; **207**(6): 990-8.
4. Shen Z, Ning F, Zhou W, et al. Superspreading SARS events, Beijing, 2003. *Emerg Infect Dis* 2004; **10**(2): 256-60.
5. Streeck H, Schulte B, Kümmerer BM, et al. Infection fatality rate of SARS-CoV2 in a super-spreading event in Germany. *Nat Commun* 2020; **11**(1): 5829.
6. Leclerc QJ, Fuller NM, Knight LE, Funk S, Knight GM. What settings have been linked to SARS-CoV-2 transmission clusters? *Wellcome Open Res* 2020; **5**: 83.
7. Corman VM, Landt O, Kaiser M, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* 2020; **25**(3).
8. Schmithausen RM, Döhla M, Schöpler H, et al. Characteristic Temporary Loss of Taste and Olfactory Senses in SARS-CoV-2-positive-Individuals with Mild Symptoms. *Pathog Immun* 2020; **5**(1): 117-20.
9. Kyriakopoulos AM, Papaefthymiou A, Georgilas N, Doulberis M, Kountouras J. The Potential Role of Super Spread Events in SARS-COV-2 Pandemic; a Narrative Review. *Arch Acad Emerg Med* 2020; **8**(1): e74.
10. Patel KP, Vunnam SR, Patel PA, et al. Transmission of SARS-CoV-2: an update of current literature. *Eur J Clin Microbiol Infect Dis* 2020; **39**(11): 2005-11.
11. Lednicky JA, Lauzardo M, Fan ZH, et al. Viable SARS-CoV-2 in the air of a hospital room with COVID-19 patients. *Int J Infect Dis* 2020; **100**: 476-82.
12. Santarpia JL, Rivera DN, Herrera VL, et al. Aerosol and surface contamination of SARS-CoV-2 observed in quarantine and isolation care. *Sci Rep* 2020; **10**(1): 12732.
13. Günther T, Czech-Sioli M, Indenbirken D, et al. SARS-CoV-2 outbreak investigation in a German meat processing plant. *EMBO Mol Med* 2020; **12**(12): e13296.
14. Pokora R, Kutschbach S, Weigl M, et al. Investigation of superspreading COVID-19 outbreak events in meat and poultry processing plants in Germany: A cross-sectional study. *PLoS One* 2021; **16**(6): e0242456.
15. Nazarenko Y. Air filtration and SARS-CoV-2. *Epidemiol Health* 2020; **42**: e2020049.
16. Vardavas CI, Nikitara K. COVID-19 and smoking: A systematic review of the evidence. *Tob Induc Dis* 2020; **18**: 20.
17. Engin AB, Engin ED, Engin A. Two important controversial risk factors in SARS-CoV-2 infection: Obesity and smoking. *Environ Toxicol Pharmacol* 2020; **78**: 103411.
18. Israel A, Feldhamer E, Lahad A, Levin-Zamir D, Lavie G. Smoking and the risk of COVID-19 in a large observational population study. *medRxiv* 2020: 2020.06.01.20118877.
19. Grundy EJ, Suddek T, Filippidis FT, Majeed A, Coronini-Cronberg S. Smoking, SARS-CoV-2 and COVID-19: A review of reviews considering implications for public health policy and practice. *Tob Induc Dis* 2020; **18**: 58.
20. Changeux JP, Amoura Z, Rey FA, Miyara M. A nicotinic hypothesis for Covid-19 with preventive and therapeutic implications. *C R Biol* 2020; **343**(1): 33-9.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

21. Paleiron N, Mayet A, Marbac V, et al. Impact of Tobacco Smoking on the Risk of COVID-19: A Large Scale Retrospective Cohort Study. *Nicotine Tob Res* 2021; **23**(8): 1398-404.

22. Li X, Xu W, Dozier M, He Y, Kirolos A, Theodoratou E. The role of children in transmission of SARS-CoV-2: A rapid review. *J Glob Health* 2020; **10**(1): 011101.

23. Li X, Xu W, Dozier M, et al. The role of children in the transmission of SARS-CoV2: updated rapid review. *J Glob Health* 2020; **10**(2): 021101.

24. Viner RM, Mytton OT, Bonell C, et al. Susceptibility to SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults: A Systematic Review and Meta-analysis. *JAMA Pediatr* 2021; **175**(2): 143-56.

Figure Legends

Fig. 1: Study participants. Of the 400 people contacted originally (left) 362 adults and 49 children agreed to enroll in the study. An overview of the number of samples collected is given on the right. Downstream sample processing included centrifugation of blood samples for plasma collection (SARS-CoV-2 ELISAs), and viral RNA extraction from swab samples (SARS-CoV-2 RT-PCR).

Fig. 2: Reconstructed 3D-Model of the venue hall. Self-administered questionnaires included questions about main seating-position of the participants during the evening event as specifying table and seat with the help of a schematic seating plan. Metric room coordinates for all tables, seats, and ventilation-points were assessed and the seating was reconstructed from pictures taken during the event. Therefore, the location of the stage, the bar, the exit as well as the tables and the air-inlets/outlets were reconstructed in a 3D-Model. The original external dimensions of the building were 27m x 13.20m x 4.20m. Tables, where more than 7 infected individuals have stayed are colored in dark red, this includes the stage and bar as well. Air-inlets are colored in violet and the air-outlets in blue. Infected participants had been seated mostly at tables close to the bar, the bar itself and on stage. One table with 8 out of 11 infected people, was located far away from the bar at the other side of the hall and close to an air inlet. The group sitting on stage showed as well high numbers of infection (18 infected out of 24). Greater proximity to air outlets seems to be associated with increased risk of infection with a crude OR=1.39 [0.86; 2.25].

Fig. 3 Odds-Ratio for the likelihood of SARS-CoV-2 infection by age groups. Participants were divided into age groups of 8, 15, or 25 years, participants younger than 18 or older than 65 years. Participants were considered to have been infected during the event if they were SARS-CoV-2 antibody positive (ELISA).

Fig. 4: Odds ratios for the association of SARS-CoV-2 infection with specific activities of the participants and their location in the venue relative to ventilation shafts. The model was additionally adjusted for age, sex, duration of attendance, participation in multiple activities, and cumulative proximity to other infected persons, and common household.

Fig. 5: Odds ratios for symptoms of SARS-CoV-2 antibody-positive participants in the 14 days following the super spreading event. The information on symptoms was derived from the self-administered questionnaire, which was filled out on the day of sample collection. Odds ratio estimates (OR) are shown with confidence intervals

Suppl. Fig. 1: Correlation of SARS-CoV-2 Euroimmun ELISA results for IgA and IgG. The correlation of IgA levels to IgG levels in the same person was significant (r: Pearson coefficient, $p < 0.0001$, 95 % CI, 0.7043 to 0.7902). The dotted lines mark the ratios above which each ELISA result is considered positive.

Suppl. Fig. 2: Correlation of plasma neutralization capacity and IgG ELISA results (Euroimmun) from each donor. The dotted line marks the ratio above which the ELISA result is considered positive. The correlation coefficient (Pearson) was

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

0.3667 (95 % CI, 0.2275 to 0.4192, $p<0.0001$). Samples with a negative result in the neutralization assay were set as 0.1 here so as to appear on the logarithmic axis.

Supplemental Figure 3: Quantile plot of observed p-values from analyses of inverse distance [1/m] to single specific study participants as risk factor for corona-virus infection. In case of no association, the ordered log-transformed p-values are expected to lie on, or below the diagonal. Panel A: results from crude analyses, Panel B: analyses were adjusted for age, sex, common household and duration of attendance.

Supplementary table 1: Estimated relative risk of SARS-CoV-2 infection (IGG-positive) from logistic regression on summary measures of spatial proximity between participants in terms of odds ratio estimates (OR) with confidence interval and p-values. a) adjusted for sex, age, common household and duration. b) multivariate analysis, mutually adjusted for distance to ventilation system, participation in (multiple) performances, going out of doors during the intermission, and participating in the grand finale. c) multivariate analysis, mutually adjusted for distance to ventilation system, participation in (multiple) performances, going out of doors during the intermission, and participating in the grand finale and adjusted for sex, age, common household and duration.




Fig. 1: Study participants. Of the 400 people contacted originally (left) 362 adults and 49 children agreed to enroll in the study. An overview of the number of samples collected is given on the right. Downstream sample processing included centrifugation of blood samples for plasma collection (SARS-CoV-2 ELISAs), and viral RNA extraction from swab samples (SARS-CoV-2 RT-PCR).

342x107mm (96 x 96 DPI)




Fig. 2: Reconstructed 3D-Model of the venue hall. Self-administered questionnaires included questions about main seating-position of the participants during the evening event as specifying table and seat with the help of a schematic seating plan. Metric room coordinates for all tables, seats, and ventilation-points were assessed and the seating was reconstructed from pictures taken during the event. Therefore, the location of the stage, the bar, the exit as well as the tables and the air-inlets/outlets were reconstructed in a 3D-Model. The original external dimensions of the building were 27m x 13.20m x 4.20m. Tables, where more than 7 infected individuals have stayed are colored in dark red, this includes the stage and bar as well. Air-inlets are colored in violet and the air-outlets in blue. Infected participants had been seated mostly at tables close to the bar, the bar itself and on stage. One table with 8 out of 11 infected people, was located far away from the bar at the other side of the hall and close to an air inlet. The group sitting on stage showed as well high numbers of infection (18 infected out of 24). Greater proximity to air outlets seems to be associated with increased risk of infection with a crude OR=1.39 [0.86; 2.25].

306x185mm (96 x 96 DPI)




Fig. 3 Odds-Ratio for the likelihood of SARS-CoV-2 infection by age groups. Participants were divided into age groups of 8, 15, or 25 years, participants younger than 18 or older than 65 years. Participants were considered to have been infected during the event if they were SARS-CoV-2 antibody positive (ELISA).

283x160mm (300 x 300 DPI)




Fig. 4: Odds ratios for the association of SARS-CoV-2 infection with specific activities of the participants and their location in the venue relative to ventilation shafts. The model was additionally adjusted for age, sex, duration of attendance, participation in multiple activities, and cumulative proximity to other infected persons, and common household.


272x124mm (300 x 300 DPI)




Fig. 5: Odds ratios for symptoms of SARS-CoV-2 antibody-positive participants in the 14 days following the super spreading event. The information on symptoms was derived from the self-administered questionnaire, which was filled out on the day of sample collection. Odds ratio estimates (OR) are shown with confidence intervals.

281x217mm (300 x 300 DPI)


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



277x218mm (300 x 300 DPI)



272x201mm (300 x 300 DPI)



337x129mm (96 x 96 DPI)

	OR	95 % confidence interval		p-value
Proximity of infected persons [sum 1/m]	0.99	0.98	1.01	0.43
Adjusted a)	1	0.98	1.02	0.96
Mutually adjusted b)	0.99	0.97	1.01	0.57
Mutually adjusted c)	0.99	0.97	1.02	0.65
Alternative consideration in distance-bands				
Infected persons within ≤ 1.5 m [count]	1.01	0.96	1.07	0.68
Infected persons in 1.5 - ≤ 3 m [count]	0.96	0.92	1	0.04
Infected persons in 3 - ≤ 4.5 m [count]	1.03	1	1.06	0.02
Infected persons within ≤ 1.5 m [count] adjusted a)	1.03	0.97	1.1	0.37
Infected persons in 1.5 - ≤ 3 m [count]	0.96	0.92	1.01	0.11
Infected persons in 3 - ≤ 4.5 m [count]	1.03	1	1.06	0.08
Infected persons within ≤ 1.5 m [count] mutually adjusted b)	1.01	0.95	1.07	0.73
Infected persons in 1.5 - ≤ 3 m [count]	0.98	0.94	1.02	0.36
Infected persons in 3 - ≤ 4.5 m [count]	1.05	1.02	1.08	0.001
Infected persons within ≤ 1.5 m [count] mutually adjusted c)	1.02	0.95	1.09	0.64
Infected persons in 1.5 - ≤ 3 m [count]	0.98	0.93	1.03	0.36
Infected persons in 3 - ≤ 4.5 m [count]	1.04	1	1.07	0.04

Supplementary table 1: Estimated relative risk of SARS-CoV-2 infection (IGG-positive) from logistic regression on summary measures of spatial proximity between participants in terms of odds ratio estimates (OR) with confidence interval and p-values. a) adjusted for sex, age, common household and duration. b) multivariate analysis, mutually adjusted for distance to ventilation system, participation in (multiple) performances, going out of doors during the intermission, and participating in the grand finale. c) multivariate analysis, mutually adjusted for distance to ventilation system, participation in (multiple) performances, going out of doors during the intermission, and participating in the grand finale and adjusted for sex, age, common household and duration.

Reporting checklist for cohort study.

Based on the STROBE cohort guidelines.

Instructions to authors

Complete this checklist by entering the page numbers from your manuscript where readers will find each of the items listed below.

Your article may not currently address all the items on the checklist. Please modify your text to include the missing information. If you are certain that an item does not apply, please write "n/a" and provide a short explanation.

Upload your completed checklist as an extra file when you submit to a journal.

In your methods section, say that you used the STROBE cohort reporting guidelines, and cite them as:

von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies.

Reporting Item			Page Number
Title and abstract			
Title	#1a	Indicate the study's design with a commonly used term in the title or the abstract	1
Abstract	#1b	Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background / rationale	#2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	#3	State specific objectives, including any prespecified hypotheses	2
Methods			
Study design	#4	Present key elements of study design early in the paper	5
Setting	#5	Describe the setting, locations, and relevant dates, including periods	7

of recruitment, exposure, follow-up, and data collection

Eligibility criteria	#6a	Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up.	5
Eligibility criteria	#6b	For matched studies, give matching criteria and number of exposed and unexposed	5
Variables	#7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5
Data sources / measurement	#8	For each variable of interest give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group. Give information separately for for exposed and unexposed groups if applicable.	7
Bias	#9	Describe any efforts to address potential sources of bias	4
Study size	#10	Explain how the study size was arrived at	5
Quantitative variables	#11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen, and why	5-6
Statistical methods	#12a	Describe all statistical methods, including those used to control for confounding	7-8
7-8			
Statistical methods	#12b	Describe any methods used to examine subgroups and interactions	
Statistical methods	#12c	Explain how missing data were addressed	
Statistical methods	#12d	If applicable, explain how loss to follow-up was addressed	
Statistical methods	#12e	Describe any sensitivity analyses	8

8

Results

Participants	#13a	Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible,	8
--------------	----------------------	---	---

		included in the study, completing follow-up, and analysed. Give information separately for for exposed and unexposed groups if applicable.	
Participants	#13b	Give reasons for non-participation at each stage	8
Participants	#13c	Consider use of a flow diagram	
8			
Descriptive data	#14a	Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders. Give information separately for exposed and unexposed groups if applicable.	8
Descriptive data	#14b	Indicate number of participants with missing data for each variable of interest	
8			
Descriptive data	#14c	Summarise follow-up time (eg, average and total amount)	
8			
Outcome data	#15	Report numbers of outcome events or summary measures over time. Give information separately for exposed and unexposed groups if applicable.	
8			
Main results	#16a	Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	8-9
Main results	#16b	Report category boundaries when continuous variables were categorized	8-9
Main results	#16c	If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
8-9			
Other analyses	#17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10-11

Discussion

1	Key results	#18	Summarise key results with reference to study objectives	12
2				
3	Limitations	#19	Discuss limitations of the study, taking into account sources of	12
4			potential bias or imprecision. Discuss both direction and magnitude of	
5			any potential bias.	
6				
7				
8	Interpretation	#20	Give a cautious overall interpretation considering objectives,	11-12
9			limitations, multiplicity of analyses, results from similar studies, and	
10			other relevant evidence.	
11				
12				
13	Generalisability	#21	Discuss the generalisability (external validity) of the study results	12
14				
15				
16	Other			
17	Information			
18				
19				
20	Funding	#22	Give the source of funding and the role of the funders for the present	15
21			study and, if applicable, for the original study on which the present	
22			article is based	
23				
24				

The STROBE checklist is distributed under the terms of the Creative Commons Attribution License CC-BY. This checklist was completed on 01. December 2021 using <https://www.goodreports.org/>, a tool made by the [EQUATOR Network](#) in collaboration with [Penelope.ai](#)

BMJ Open

Dynamics, outcomes, and prerequisites of the first SARS-CoV-2 superspreading event in Germany, in February 2020: a cross-sectional epidemiological study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-059809.R1
Article Type:	Original research
Date Submitted by the Author:	23-Feb-2022
Complete List of Authors:	Wessendorf, Lukas; University Hospital Bonn, Virology Richter, Enrico; University Hospital Bonn, Virology Schulte, Bianca; University Hospital Bonn, Institute for Virology Schmithausen, Ricarda Maria; University Hospital Bonn, Hygiene Exner, Martin; University Hospital Bonn Lehmann, Nils; University of Duisburg-Essen, Institute for Medical Informatics, Biometry and Epidemiology Coenen, Martin; Rheinische Friedrich Wilhelms Universität Bonn, Clinical Study Core Unit, Study Center Bonn (SZB) Fuhrmann, Christine; University Hospital Bonn Kellings, Angelika; University Hospital Bonn Hüsing, Anika; University of Duisburg-Essen, Institute of Medical Informatics, Biometry and Epidemiology Jöckel, Karl-Heinz; University of Duisburg-Essen, Institute of Medical Informatics, Biometry and Epidemiology Streeck, Hendrik; German Centre for Infection Research
Primary Subject Heading:	Infectious diseases
Secondary Subject Heading:	Public health
Keywords:	COVID-19, VIROLOGY, Infection control < INFECTIOUS DISEASES

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Dynamics, outcomes, and prerequisites of the first SARS-CoV-2 superspreading event in Germany, in February 2020: a cross-sectional epidemiological study

Lukas Wessendorf¹, Enrico Richter¹, Bianca Schulte¹, Ricarda Maria Schmithausen², Martin Exner², Nils Lehmann³, Martin Coenen⁴, Christine Fuhrmann⁴, Angelika Kellings⁴, Anika Hüsing³, Karl-Heinz Jöckel³, and Hendrik Streeck^{1,*}

¹ Institute of Virology, University Hospital, University of Bonn, Germany, and German Center for Infection Research (DZIF), partner site Bonn-Cologne

² Institute of Hygiene and Public Health, University Hospital, University of Bonn, Germany

³ Institute of Medical Informatics, Biometry und Epidemiology (IMIBE), University Hospital Essen, Germany

⁴ Clinical Study Core Unit, Study Center Bonn (SZB), Institute of Clinical Chemistry and Clinical Pharmacology, University Hospital Bonn, Germany

* Corresponding author: Hendrik Streeck, Institute of Virology, University Hospital Bonn, Venusberg-Campus 1, 53127 Bonn, Germany, Hendrik.Streeck@ukbonn.de

Word Count: 4,581

Keywords: SARS-CoV-2, COVID-19, superspreading, cross-sectional study, ventilation

Abstract

Objectives

The first German SARS-CoV-2 outbreak was a superspreading event in Gangelt, North Rhine-Westphalia during indoor carnival festivities called “Kappensitzung” (15th of February 2020). We determined SARS-CoV-2 RT-PCR positivity rate, SARS-CoV-2-specific antibodies, and analyzed the conditions and dynamics of superspreading, including ventilation, setting dimensions, distance from infected persons and behavioral patterns.

Design

In a cross-sectional epidemiological study (51 days post-event), participants were asked to give blood, pharyngeal swabs and complete self-administered questionnaires.

Setting

1
2
3 42 The SARS-CoV-2 superspreading event took place during festivities in the small
4
5 43 community of Gangelt in February 2020. This 5 h event included 450 people (6-79
6
7 44 years of age) in a building of 27m x 13.20m x 4.20m.

8 45 **Participants**

9
10 46 Out of 450 event participants, 411 volunteered to participate in this study.

11
12 47 **Primary and Secondary Outcome Measures**

13 48 Primary outcome: infection status (determined by IgG ELISA). Secondary outcome:
14
15 49 symptoms (determined by questionnaire).

16
17 50 **Results**

18
19 51 Overall, 46% (n=186/404) of participants had been infected, and their spatial
20
21 52 distribution was associated with proximity to the ventilation system (OR 1.39, 95% KI
22
23 53 [0.86; 2.25]). Risk of infection was highly associated with age: children (OR: 0.33
24
25 54 [0.267; 0.414]) and young adults (age 18-25) had a lower risk of infection than older
26
27 55 participants (average risk increase of 28% per 10 year). Behavioral differences were
28
29 56 also risk-associated including time spent outside (OR: 0.55 [0.33; 0.91]) or smoking
30
31 57 (OR: 0.32 [0.124; 0.81]).

32
33 58 **Conclusions**

34
35 59 Our findings underline the importance of proper indoor ventilation for future events.
36
37 60 Lower susceptibility of children/young adults indicates their limited involvement in
38
39 61 superspreading.

40
41 62

42
43 63

44
45 64 **Strengths and limitations of this study**

46
47 65

48
49 66

50
51 67

52
53 68

54
55 69

56
57 70

58
59 71

60
61 72

62
63 73

64
65 74

66
67 75

68
69 76

70
71 77

72
73 78

74
75 79

76
77 80

78
79 81

80
81 82

82
83 83

84
85 84

86
87 85

88
89 86

90
91 87

92
93 88

94
95 89

96
97 90

98
99 91

100
101 92

102
103 93

104
105 94

106
107 95

108
109 96

110
111 97

112
113 98

114
115 99

116
117 100

118
119 101

120
121 102

122
123 103

124
125 104

126
127 105

128
129 106

130
131 107

132
133 108

134
135 109

136
137 110

138
139 111

140
141 112

142
143 113

144
145 114

146
147 115

148
149 116

150
151 117

152
153 118

154
155 119

156
157 120

158
159 121

160
161 122

162
163 123

164
165 124

166
167 125

168
169 126

170
171 127

172
173 128

174
175 129

176
177 130

178
179 131

180
181 132

182
183 133

184
185 134

186
187 135

188
189 136

190
191 137

192
193 138

194
195 139

196
197 140

198
199 141

200
201 142

202
203 143

204
205 144

206
207 145

208
209 146

210
211 147

212
213 148

214
215 149

216
217 150

218
219 151

220
221 152

222
223 153

224
225 154

226
227 155

228
229 156

230
231 157

232
233 158

234
235 159

236
237 160

238
239 161

240
241 162

242
243 163

244
245 164

246
247 165

248
249 166

250
251 167

252
253 168

254
255 169

256
257 170

258
259 171

260
261 172

262
263 173

264
265 174

266
267 175

268
269 176

270
271 177

272
273 178

274
275 179

276
277 180

278
279 181

280
281 182

282
283 183

284
285 184

286
287 185

288
289 186

290
291 187

292
293 188

294
295 189

296
297 190

298
299 191

300
301 192

302
303 193

304
305 194

306
307 195

308
309 196

310
311 197

312
313 198

314
315 199

316
317 200

318
319 201

320
321 202

322
323 203

324
325 204

326
327 205

328
329 206

330
331 207

332
333 208

334
335 209

336
337 210

338
339 211

340
341 212

342
343 213

344
345 214

346
347 215

348
349 216

350
351 217

352
353 218

354
355 219

356
357 220

358
359 221

360
361 222

362
363 223

364
365 224

366
367 225

368
369 226

370
371 227

372
373 228

374
375 229

376
377 230

378
379 231

380
381 232

382
383 233

384
385 234

386
387 235

388
389 236

390
391 237

392
393 238

394
395 239

396
397 240

398
399 241

400
401 242

402
403 243

404
405 244

406
407 245

408
409 246

410
411 247

412
413 248

414
415 249

416
417 250

418
419 251

420
421 252

422
423 253

424
425 254

426
427 255

428
429 256

430
431 257

432
433 258

434
435 259

436
437 260

438
439 261

440
441 262

442
443 263

444
445 264

446
447 265

448
449 266

450
451 267

452
453 268

454
455 269

456
457 270

458
459 271

460
461 272

462
463 273

464
465 274

466
467 275

468
469 276

470
471 277

472
473 278

474
475 279

476
477 280

478
479 281

480
481 282

482
483 283

484
485 284

486
487 285

488
489 286

490
491 287

492
493 288

494
495 289

496
497 290

498
499 291

500
501 292

502
503 293

504
505 294

506
507 295

508
509 296

510
511 297

512
513 298

514
515 299

516
517 300

518
519 301

520
521 302

522
523 303

524
525 304

526
527 305

528
529 306

530
531 307

532
533 308

534
535 309

536
537 310

538
539 311

540
541 312

542
543 313

544
545 314

546
547 315

548
549 316

550
551 317

552
553 318

554
555 319

556
557 320

558
559 321

560
561 322

562
563 323

564
565 324

566
567 325

568
569 326

570
571 327

572
573 328

574
575 329

576
577 330

578
579 331

580
581 332

582
583 333

584
585 334

586
587 335

588
589 336

590
591 337

592
593 338

594
595 339

596
597 340

598
599 341

600
601 342

602
603 343

604
605 344

606
607 345

608
609 346

610
611 347

612
613 348

614
615 349

616
617 350

618
619 351

620
621 352

622
623 353

624
625 354

626
627 355

628
629 356

630
631 357

632
633 358

634
635 359

636
637 360

638
639 361

640
641 362

642
643 363

644
645 364

646
647 365

648
649 366

650
651 367

652
653 368

654
655 369

656
657 370

658
659 371

660
661 372

662
663 373

664
665 374

666
667 375

668
669 376

670
671 377

672
673 378

674
675 379

676
677 380

678
679 381

680
681 382

682
683 383

684
685 384

686
687 385

688
689 386

690
691 387

692
693 388

694
695 389

696
697 390

698
699 391

700
701 392

702
703 393

704
705 394

706
707 395

708
709 396

710
711 397

712
713 398

714
715 399

716
717 400

718
719 401

720
721 402

722
723 403

724
725 404

726
727 405

728
729 406

730
731 407

732
733 408

734
735 409

736
737 410

738
739 411

740
741 412

742
743 413

744
745 414

746
747 415

748
749 416

750
751 417

752
753 418

754
755 419

756
757 420

758
759 421

760
761 422

762
763 423

764
765 424

766
767 425

768
769 426

770
771 427

772
773 428

774
775 429

776
777 430

778
779 431

780
781 432

782
783 433

784
785 434

786
787 435

788
789 436

790
791 437

792
793 438

794
795 439

796
797 440

798
799 441

800
801 442

802
803 443

804
805 444

806
807 445

808
809 446

810
811 447

812
813 448

814
815 449

816
817 450

818
819 451

820
821 452

822
823 453

824
825 454

826
827 455

828
829 456

830
831 457

832
833 458

834
835 459

836
837 460

838
839 461

840
841 462

842
843 463

844
845 464

846
847 465

848
849 466

850
851 467

852
853 468

854
855 469

856
857 470

858
859 471

860
861 472

862
863 473

864
865 474

866
867 475

868
869 476

870
871 477

872
873 478

874
875 479

876
877 480

878
879 481

880
881 482

882
883 483

884
885 484

886
887 485

888
889 486

890
891 487

892
893 488

894
895 489

896
897 490

898
899 491

900
901 492

902
903 493

904
905 494

906
907 495

908
909 496

910
911 497

912
913 498

914
915 499

916
917 500

918
919 501

920
921 502

922
923 503

924
925 504

926
927 505

928
929 506

930
931 507

932
933 508

934
935 509

936
937 510

938
939 511

940
941 512

942
943 513

944
945 514

946
947 515

948
949 516

950
951 517

952
953 518

954
955 519

956
957 520

958
959 521

960
961 522

962
963 523

964
965 524

966
967 525

968
969 526

970
971 527

972
973 528

974
975 529

976
977 530

978
979 531

980
981 532

982
983 533

984
985 534

986
987 535

988
989 536

990
991 537

992
993 538

994
995 539

996
997 540

998
999 541

1000
1001 542

1002
1003 543

1004
1005 544

1006
1007 545

1008
1009 546

1010
1011 547

1012
1013 548

1014
1015 549

1016
1017 550

1018
1019 551

1020
1021 552

1022
1023 553

1024
1025 554

1026
1027 555

1028
1029 556

1030
1031 557

1032
1033 558

1034
1035 559

1036
1037 560

1038
1039 561

1040
1041 562

1042
1043 563

1044
1045 564

1046
1047 565

1048
1049 566

1050
1051 567

1052
1053 568

1054
1055 569

1056
1057 570

1058
1059 571

1060
1061 572

1062
1063 573

1064
1065 574

1066
1067 575

1068
1069 576

1070
1071 577

1072
1073 578

1074
1075 579

1076
1077 580

1078
1079 581

1080
1081 582

1082
1083 583

1084
1085 584

1086
1087 585

1088
1089 586

1090
1091 587

1092
1093 588

1094
1095 589

1096
1097 590

1098
1099 591

1100
1101 592

1102
1103 593

1104
1105 594

1106
1107 595

1108
1109 596

1110
1111 597

1112
1113 598

1114
1115 599

1116
1117 600

1118
1119 601

1120
1121 602

1122
1123 603

1124
1125 604

1126
1127 605

1128
1129 606

1130
1131 607

1132
1133 608

1134
1135 609

1136
1137 610

1138
1139 611

1140
1141 612

1142
1143 613

1144
1145 614

1146
1147 615

1148
1149 616

1150
1151 617

1152
1153 618

1154
1155 619

1156
1157 620

1158
1159 621

1160
1161 622

1162
1163 623

1164
1165 624

1166
1167 625

1168
1169 626

1170
1171 627

1172
1173 628

1174
1175 629

1176
1177 630

1178
1179 631

1180
1181 632

1182
1183 633

1184
1185 634

1186
1187 635

1188
1189 636

1190
1191 637

1192
1193 638

1194
1195 639

1196
1197 640

1198
1199 641

1200
1201 642

1202
1203 643

1204
1205 644

1206
1207 645

1208
1209 646

1210
1211 647

1212
1213 648

1214
1215 649

1216
1217 650

1218
1219 651

1220
1221 652

1222
1223 653

1224
1225 654

1226
1227 655

1228
1229 656

1230
1231 657

1232
1233 658

1234
1235 659

1236
1237 660

1238
1239 661

1240
1241 662

1242
1243 663

1244
1245 664

1246
1247 665

1248
1249 666

1250
1251 667

1252
1253 668

1254
1255 669

1256
1257 670

1258
1259 671

1260
1261 672

1262
1263 673

1264
1265 674

1266
1267 675

1268
1269 676

1270
1271 677

1272
1273 678

1274
1275 679

1276
1277 680

1278
1279 681

1280
1281 682

1282
1283 683

1284
1285 684

1286
1287 685

1288
1289 686

1290
1291 687

1292
1293 688

1294
1295 689

1296
1297 690

1298
1299 691

1300
1301 692

1302
1303 693

1304
1305 694

1306
1307 695

1308
1309 696

1310
1311 697

1312
1313 698

1314
1315 699

1316
1317 700

1318
1319 701

1320
1321 702

1322
1323 703

1324
1325 704

1326
1327 705

1328
1329 706

1330
1331 707

1332
1333 708

1334
1335 709

1336
1337 710

1338
1339 711

1340
1341 712

1342
1343 713

1344
1345 714

1346
1347 715

1348
1349 716

1350
1351 717

1352
1353 718

1354
1355 719

1356
1357 720

1358
1359 721

1360
1361 722

1362
1363 723

1364
1365 724

1366
1367 725

1368
1369 726

1370
1371 727

1372
1373 728

1374
1375 729

1376
1377 730

1378
1379 731

1380
1381 732

1382
1383 733

138

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a highly transmissible and pathogenic RNA virus that emerged in late 2019 and has caused a pandemic threatening human health and public safety worldwide.¹ While factors shaping the dynamics of a pandemic are multifactorial, virulence and reproductive number are important properties of a virus.² For SARS-CoV-2 there is a substantial over-dispersion of the secondary infection distribution (individual R_0) for an individual infected with SARS-CoV-2.² An over-dispersed R_0 means that most infected people do not transmit (individual $R_0 = 0$) while a minority of infected people are super-spreaders (individual $R_0 > 5$). Superspreading has been observed for many infectious pathogens, such as measles or SARS.³ During the SARS pandemic in 2003 a superspreading event was defined as one infected person infecting eight others.⁴ For SARS-CoV-2 it has been estimated that 80% of the infections are caused by 10% of infected individuals highlighting the importance of the cluster factor (k).² In Germany an indoor carnival event in the beginning of 2020 is considered as the first major outbreak in a German city and was considered a hotspot during the beginning of the pandemic in Germany.⁵ Other SARS-CoV-2 superspreading events worldwide have been linked to indoor gatherings with close proximity of individuals.⁶ Nevertheless, most of the reported superspreading events had less than 100 cases and the reports are limited by missing data or a reporting bias.⁶

Here, we closely examined the prerequisite of a unique super-spreading event in Germany during the SARS-CoV-2 pandemic, where nearly half of the participants became infected including children. We systematically analyzed infection rate, potential individual, and environmental risk factors for infection as well as the role of the ventilation system.

Materials and methods

Study design and sampling

This cross-sectional epidemiological study was conducted 51 days after a carnival celebration in the beginning of 2020. Eleven days after the event authorities sent all known participants into quarantine after testing 38 out of 99 individuals PCR-positive.

1
2
3 108 All adults known to have attended the event were invited to participate in the study.
4
5 109 About 450 persons attended the event of which 411 participated in the study (**figure**
6
7 110 **1**, participation rate 91.3%). All study participants provided written informed consent
8
9 111 before enrolment. The Ethics Committee of the Medical Faculty of the University of
10
11 112 Bonn approved the study (approval number 085/20).
12
13 113 Self-administered questionnaires included questions about demographic background,
14
15 114 symptoms of viral infection as well as detailed information about the behavior during
16
17 115 the event. Participants' arrival and exit times were assessed in 1-hour categories.
18
19 116 Study participants were asked to provide blood specimens and pharyngeal swabs for
20
21 117 further analysis. The local health department supplied data on hospitalizations and
22
23 118 fatalities in our cohort (manuscript submitted elsewhere).
24
25 119

26 120 **Patient and public involvement**

27 121 This study was designed in close collaboration with both the local health department
28
29 122 of Heinsberg and the 'Council of 11' of Gangelt, the organizers of the event described
30
31 123 herein. The organizers as well as the city's head councilman were also involved in
32
33 124 recruitment by appealing to the local population to participate in the study. Since the
34
35 125 community of Gangelt was the center of the first German outbreak of SARS-CoV-2,
36
37 126 there was a great interest from the local public to participate in this study to help
38
39 127 understand this new virus and to gain access to early testing. Accordingly, the Ministry
40
41 128 of Labor, Health, and Social Affairs of the state government funded this study. In turn,
42
43 129 as a service to the public we informed each participant of their PCR and ELISA result
44
45 130 via letter and offered a phone hotline for questions about the results.
46
47 131

48 132 **Spatial information and description of the event**

49 133 The event took place on February 15th, 2020 and consisted of speeches, dance, and
50
51 134 music performances for a total of five hours, with one large intermission. It was a
52
53 135 ticketed event, where ticket sale was open to the public. Most of the participants were
54
55 136 inhabitants of Gangelt. It was hosted at a small community center (320 square meters)
56
57 137 in a single open space with a bar in the front close to the entrance and a stage at the
58
59 138 back. The tables, each with two benches, were arranged in two blocks with a center
60
139 aisle towards the stage. Alcoholic and nonalcoholic drinks were served in glasses and
140
141 a food truck was located outside in front of the venue. While most participants (about
450 people, 1.4 individuals per square meter) were sitting in the hall, a committee of

eleven individuals hosting the event were sitting on stage. The eleven people on stage switched after a break.

Metric room coordinates (length and width [m]) for areas, tables, benches and ventilation shafts were assessed via measurements, seating plan and photos from the event. Persons providing multiple positions were considered as spending an equal amount of time on different positions. When exact seating was unclear and information was available on table or greater area localisation (bar, stage), average coordinate values were used. On the grounds of these coordinates, we calculated pairwise metric distances between all persons and distances to closest inlet and outlet airshafts. For all persons their pairwise inverse distances were summarized as mean inverse distance. Inverse metric distances to persons or airshafts were regarded as representing infectious potential through local proximity, and inverse distances were capped at 2.5 (the inverse of the width of a seat of 0.4 m). Alternatively, we counted all infected persons within adjacent rings of 1.5 m width around each participant as a measure of crowdedness and infectious potential.

Pharyngeal swab and blood preparation

Pharyngeal swabs of participants were performed with FLOQSwabs (Copan) and immediately stored in UTM RT-mini tubes containing UTM Viral Stabilization Media (Copan) at 4 °C. Venous blood was drawn into EDTA tubes (Sarstedt) per participant and was transported to the laboratory at the University Hospital Bonn.

Anti-SARS-CoV-2 ELISA

Anti-SARS-CoV-2 IgA and IgGs were determined using enzyme-linked immunosorbent assays (ELISA) on the EUROIMMUN Analyzer I platform (EI 2606-9601 A, and EI2606-9601 G, respectively).⁵ A result was considered positive when a ratio (extinction of sample/extinction of calibrator) of 0.8 or higher was reached. The guidelines of the German Medical Association (RiliBÄK) were abided by, including internal and external quality controls.

Reverse transcription polymerase chain reaction (RT-PCR)

Viral RNA was extracted from each 300µl swab sample via the chemagic Viral 300 assay (according to manufacturer's instructions) on the Perkin Elmer chemagic™ Prime™ instrument platform. The presence of two viral target genes (E and RdRP) was assessed in each sample by real time RT-PCR (SuperScript™III One-Step RT-

PCR System with Platinum™ TaqDNA Polymerase, Thermo Fisher). The following primers were used, for E gene: E_Sarbeco_F1 and R, and probe E_Sarbeco_P1, for RdRP gene: RdRP_SARSr_F, and R, and probe RdRP_SARSr-P2.⁷ In addition, an internal control for RNA extraction, reverse transcription, and amplification was applied to each sample (innuDETECT Internal Control RNA Assay, Analytik Jena #845-ID-0007100). If amplification occurred in both virus-specific reactions samples were considered positive.

SARS-CoV-2 neutralization assay

A plaque reduction neutralization test was used to determine SARS-CoV-2 neutralization capacity as previously described.⁵ Briefly, plasma samples were heat-inactivated and supernatant transferred to a new tube and serially two-fold diluted in OptiPRO™SFM (Gibco). 120 µL of each plasma dilution was mixed with 80 plaque-forming units (PFU) of SARS-CoV-2 in 120 µL OptiPRO™SFM cell culture medium and used to infect Vero E6 cells (1.25x10⁵ cells/well seeded into 24-well plates 24 h before). Subsequently, the inoculum was removed and cells were overlaid with a mixture of carboxymethylcellulose (Sigma) and 2xMEM (Biochrom). Following 3-day incubation, the overlay was removed and the 24-well plates were fixed using a 6% (v/v) formaldehyde solution and stained with 1% (w/v) crystal violet in 20% ethanol revealing the formation of plaques. Finally, the neutralizing titers were calculated as the reciprocal of serum dilutions resulting in neutralization of 50% input virus (NT50), read out as reduction in the number of plaques.

Data management and quality control

The Clinical Study Core Unit of the Study Center Bonn (SZB) supported the study by outlining the study protocol and developing the informed consent form as well as participants information sheets with respect to data management and quality control. The data were gathered on paper-based Case Report Forms (pCRF). Data was entered as double-data-entry into the REDCap study database programmed and hosted by SZB. Study personnel was trained by experienced members of the SZB. A quality manager was on site to support the study team. Monitoring of trial data and informed consent forms was performed according to the monitoring plan by qualified SZB staff. The ethics committee of the Medical Faculty of the University of Bonn was involved and approved the study (reference no. 085/20)

Statistical analysis

Associations between positive infection status (defined as an IgG ratio ≥ 0.8), and exposure variables were analysed via logistic regression models. Exposure variables were included crudely, and adjusted for the potential confounding factors age, sex, and duration of attendance as fixed effects. To correct for common household effects a random effects model was used. We present odds ratios with 95% confidence intervals. Because we present data on a single specific event among a limited number of participants, we completely refrain from presenting p-values. All analyses were done with SAS 9.4.

Results

411 out of an estimated 450 participants of the event responded to our study invitation, resulting in a response rate of 91.3%. 404 individuals provided plasma samples and 316 pharyngeal swabs (**figure 1**). Genders were represented equally among all 404 participants (n= 201/404, 50% were male) with a broad range in age (6-79 years, median age 36 years) and level of education (**table 1**). 297 individuals were residents of the community the event took place in, 103 lived in other parts of the county, and 11 were external visitors. In total five participants of the event were hospitalized and one participant subsequently died.

Overall, 186 out of 404 individuals tested seropositive for IgG- and 161 for IgA-antibodies (**suppl. figure 1**). To confirm seropositivity we performed a plaque reduction neutralization assay (**suppl. figure 2**) demonstrating neutralizing activity against SARS-CoV-2 of their respective antibody responses. Given the low specificity of the IgA assay, IgA seropositivity was not further considered.⁵ 19 participants tested positive in RT-PCR; these were considered infected during the superspreading event only if they were also IgG-positive (this was the case with 16 out of the 19 participants).

Overall, we found that (n= 186/404) 46.0% (95% CI: [41.2%; 51.0%]) tested seropositive who attended the event, which was significantly higher than the overall estimated infection rate in the same community at large at that time. Indeed, officially 3.1% of the community were reported as positive cases at that time, but we estimated the infection rate to be 15.5% (95% CI:[12.3%; 19.0%])⁵ for the community. Taken

1
2
3 244 together, an estimated 46% of participants became infected during a single
4
5 245 superspreading event.
6
7 246
8 247 No association between the gender of participants and risk of infection was found ((OR:
9
10 248 1.01 [0.65; 1.58]) for women). On average infected individuals had a higher body mass
11 249 index (26.2kg/m² compared to 24.3kg/m² for uninfected individuals). Infected
12
13 250 participants were more likely to be clustered living in the same household (**table 1**).
14
15 251 Having at least one comorbidity, including lung disease (n= 11/26, 42.3%),
16 252 cardiovascular disease (n= 8/15, 53.3%), neurological disease (n= 1/6, 16.7%), cancer
17
18 253 (n= 7/12) (58.3%) or diabetes (n= 4/5, 80%), did not increase the risk of infection (OR:
19
20 254 0.64 [0.33; 1.26]). We next assessed whether age influenced the risk of infection at the
21
22 255 event, considering gender, duration of attendance and common household as
23
24 256 covariates. Comparison across age-categories showed a lower risk for children (OR:
25 257 0.31 [0.14; 0.69]), and also for young adults (18-25 years, OR: 0.53 [0.26; 1.09]) as
26
27 258 well as adults between 25 and 40 years (OR: 0.48 [0.28; 0.85]) in comparison to older
28
29 259 adults (40 to 65 years) (OR: 1, reference), while seniors had a slightly higher risk (older
30 260 than 65 years, OR: 1.1 [0.31; 3.97]) (**figure 2**). Our data suggests that an additional 10
31
32 261 years of age were on average associated with 28% increased risk of infection (OR:
33
34 262 1.28 [1.10; 1.48]).
35
36 263 To understand the spreading dynamics of SARS-CoV-2 during the event, we first
37 264 performed a detailed analysis of potential risk factors and social behavior. We first
38
39 265 analyzed whether the ventilation system influenced the distribution of SARS-CoV-2
40
41 266 infected individuals. It is important to state that the system's air flow consisted of 75%
42 267 used and 25% fresh air. The air flow can be described as clockwise. The air system
43
44 268 uses vents along one side of the venue and on stage to take in air (**figure 3**, air inlets
45 269 purple). After 25% of fresh air has been added and the air has been filtered, vents
46
47 270 along the other side of the venue return the air into the room (**figure 3**, air outlets blue).
48
49 271 All ventilation points received the same amount of air due to throttle valves. For noise
50
51 272 protection reasons windows remained closed. The air-system used F7-Filters (ISO
52
53 273 ePM $\geq 2,5$) and had an air volume flow of 7500 m³/h.
54
55 274
56 275 Most tables located close to the air-inlets and showed no or only few infections (**figure**
57
58 276 **3**, green) also most surrounding tables showed low numbers of infection (**figure 3**,
59
60 277 yellow). Tables close to the air-outlets show high (4 or 5 infected per table) and very
278 high (6 or 7 infected per table) numbers of infected individuals. Infected participants

had been seated mostly at tables close to the bar, at the bar, or on stage. One table with 8 out of 11 infected people, was located far away from the bar at the other side of the hall and close to an air inlet. The group sitting on stage showed high numbers of infection (18 infected out of 24, **table 1**). Of note is that the overall number of participants per table was not equal for all tables. Greater proximity to air outlets was associated with increased risk of infection with a crude OR=1.39 [0.86; 2.25]. This association remained stable and was hardly attenuated from adjustment for proximity to air inlet, age, gender, duration of attendance, proximity to other infected persons, stage-activity and going outside during the intermission (**figure 4**, multiple adjusted OR=1.26 [0.63; 2.50]). A similar apparent effect for proximity to air inlets (crude OR=1.17 [0.72; 1.89]) disappeared when duration of attendance was added to the model (**figure 4**, multiple adjusted OR=1.01 [0.53; 1.94]). Overall, however, we found the increased risk for individuals located closer to the air outlet remarkably persistent (**figure 4**).

We further studied the sum of the inverse distance to all infected participants as a measure of proximity to either one common virus source or mutual infection. However, there was no evidence for increased risk of infection from greater proximity to other infected persons (**suppl. table 1**). Furthermore, we found no evidence for a single person being the source of the infection from the quantile-plot of p-values from 401 analyses conducted separately for each participant as potential source of infection (**suppl. figure 3**).

To understand the association of risk with behavior patterns we next investigated the influence of several factors on SARS-CoV-2 infection including time spent outside, smoking, performing on stage and participation during the final act ("Finale") for 30 minutes. Results were all adjusted for age, sex, common household, and duration of attendance. Participation in multiple performances was associated with slightly increased risk of infection (OR per performance: 1.08 [0.91; 1.27]), results for participation in the last "Finale" were stronger (OR: 1.41 [0.65; 3.02]), although neither was significant (**figure 4**). Duration of attendance was persistently and strongly associated with an increased infection risk of 32% with each additional hour spent at the party (OR per hour: 1.32 [1.16; 1.49]). All other analyses were adjusted for this variable as potential confounding factor.

313

314 We next determined the level of alcohol consumption as number of drinks (high-proof

315 liquor or beer) and did not observe any influence for the amount of alcohol consumption

316 on the risk of becoming infected (OR per drink: 1.00 [0.96; 1.05]). Furthermore,

317 participants who spent the break outside were less likely to be infected (OR: 0.55 [0.33;

318 0.91]) compared to individuals who spent the break inside the venue hall (**figure 4**).

319 Interestingly, however, when we determined the impact of being regular smoker

320 (defined as smoking of at least 10 cigarettes a day) on the risk of SARS-CoV-2 infection

321 we observed a reduced risk of infection (OR: 0.32 [0.12; 0.81]) even after adjustment

322 for “time spent outside”. Taken together, our results demonstrated that the duration of

323 attendance at the carnival party correlated with an increased risk of infection, but the

324 number of alcoholic drinks was not associated with infection risk, while regular smoking

325 and spending the break of the event outside showed a negative correlation with the

326 risk of infection.

327

328 We next stratified seropositive individuals by their reported symptoms. Odds-ratios for

329 each symptom were calculated for the timespan of 14 days following the event (**figure**

330 **5**). We identified that loss of smell (OR: 8.78 [4.81; 16.02]) and taste (OR: 10.09 [5.13;

331 19.88]) exhibited the strongest association with SARS-CoV-2 infection. Other

332 symptoms which were strongly associated with COVID-19 were: sweats and chills

333 (OR: 5.28 [3.08; 9.07]), muscle and joint ache (OR: 5.19 [3.19; 8.44]), fatigue (OR:

334 4.22 [2.76; 6.45]) and fever (OR: 3.73 [2.10; 6.63]) (**figure 5**). Importantly, 15.1%

335 (28/186) of the infected individuals reported no symptoms at all in a period of 14 days

336 after the event. The rate of asymptomatic infections of participants of the event was

337 lower than generally observed in the community where the event took place (36%).⁵

338 Overall, there was a lower proportion of asymptomatic cases among individuals

339 infected after the event compared to members of the community, while loss of smell

340 and taste showed the strongest association with an infection.

341

342 **Discussion**

343 The high overdispersion characteristics of SARS-CoV2 and its ability to be transmitted

344 via aerosols under certain conditions are one of the main reasons that the beginning

345 of the SARS-CoV2 pandemic was shaped by superspreading events.^{8,9} Germany`s

first superspreading event was an indoor carnival event in the beginning of 2020 in a rural community. In this naturally occurring experiment, we found that nearly half of the participants became infected and determined multiple prerequisites for superspreading and risk factors for becoming infected. While our study population is not a representative sample of the general population the event may be regarded as exemplary for similar party occasions and may help reduce the number of those infected in the future. At the time of the event described herein SARS-CoV-2 had not diversified yet, but ever since many variants of the virus have arisen and have taken turns dominating the global pandemic. Therefore, the results shown here need to be viewed as qualified in describing a superspreading event under the circumstances in the beginning of the pandemic. However, they help us to understand infection dynamics and requisites for infection with this virus family, ultimately giving a frame of reference for similar studies conducted throughout the alpha, delta, and omicron waves of the COVID-19 pandemic.

An important factor associated with infection risk was the ventilation system and the individual proximity to the ventilation outlets. Individuals close to the air-outlets that contained air with low amount of fresh air had the highest infection risk compared to those close to the air-inlets. This was particularly interesting, because we did not see any increased risk of infection from greater proximity to other infected persons, which indicates that ventilation was perhaps more important than physical proximity. Our findings are in line with previous studies that demonstrated SARS-CoV-2 to be able to become air-borne under certain conditions and that the ventilation system can have an influence on virus spread.^{10,11,12} The air filters in the venue were not capable of intercepting virus particles supporting the notion on the importance of proper indoor ventilation systems.^{13,14} Indeed, spending the break of the event outside decreased the possibility of infection underscoring the benefit of proper ventilation or fresh air to lower the amount of aerosols. Due to the nature of the event, the spatial distribution of the participants was not fixed throughout the evening, and not perfectly recapitulated, so this information carries some error. However, allowing for multiple positions per person we used all available information. Assuming further error in the spatial data to be random, this might lead to a dilution of effects, i.e. true associations may remain undetected. Complementary analyses including e.g. the persons' functions during the event show consistent results, so we see no evidence suggesting bias in our findings.

1
2
3 380 Nevertheless, the infection-rate might be overestimated as the study was conducted
4
5 381 51 days after the event as participants could have become infected not related to the
6
7 382 event. However, this weakness is limited by the official shut down of the community
8
9 383 shortly after the event: A detailed timeline of the containment measures put in place
10
11 384 after the superspreading event is included in Streeck et al.⁵. Briefly, a strict home
12
13 385 quarantine for all attendees of the carnival event was imposed after 38 out of 99
14
15 386 participants tested positive for SARS-CoV-2. In addition, 13 days after the event the
16
17 387 town went into full lockdown, including the closing of schools, childcare and outpatient
18
19 388 care facilities, and restrictions of public access to the town. These concerted
20
21 389 containment measures proved so effective that the peak of new infections in the
22
23 390 community was already reached 27 days after the event.

24 391
25 392 The consumption of alcoholic drinks did not increase the risk of infection. While it has
26
27 393 been assumed that the alcoholic effect of decreased social inhibition may increase
28
29 394 likelihood of infection, we did not find any evidence for this association questioning
30
31 395 measures of a ban on alcohol to reduce numbers of infected. It is known that current
32
33 396 and former smokers disproportionately suffer from severe COVID-19 and their
34
35 397 numbers are relatively increased among those patients that need intensive care
36
37 398 treatment compared to non-smokers^{15,16}. However, it has been previously speculated
38
39 399 that the risk of infection is lower for smokers.¹⁷ Furthermore, a meta-analysis of seven
40
41 400 studies suggests that smokers have a reduced risk of testing positive for SARS-CoV-
42
43 401 2.¹⁸ Interestingly, we also observed that regular smoking lowered the risk of infection.
44
45 402 The association might for example be explained by a role of the nicotinic acetylcholine
46
47 403 receptor (nAChR).¹⁹ Because other viruses, such as rabies virus, have been known to
48
49 404 bind nAChRs, it was hypothesized recently, that SARS-CoV-2 spike protein might bind
50
51 405 nAChRs as a coreceptor for infection.^{20,21} Indeed, *in silico* molecular docking
52
53 406 simulations predicted binding of spike to nAChRs.²² If this interaction proves to be of
54
55 407 advantage to the virus, then nicotine or its derivatives which bind nAChRs could
56
57 408 compete with SARS-CoV-2 for binding and thereby reduce interactions of the virus
58
59 409 with its target cells. Currently, at least one prospective observational study is being
60
61 410 undertaken on the effects of smoking on COVID-19 infection rates, including a smoking
62
63 411 cessation control group on nicotine substitutes.²³ While we strongly advise that
64
65 412 smoking should not be considered as a protective habit to prevent risk of infection, this

knowledge may lead to the investigation of a therapeutic or prophylactic treatment on the basis of this molecular target.²⁴

Our results indicate a trend that younger people are less likely to be infected compared to older age groups. This trend is strongest for people under 18 but levels out over 40 years of age. The risk of infection for children in superspreading events has not been investigated but the overall risk for infection in children seems to be lower than for adults as a systematic review and its recent update reported, which is further supported by our findings.^{25,26} Considering the risk of infection with SARS-CoV-2 in general however, in a meta-analysis Madewell et al. conclude that the secondary attack rate in households is lower to children contacts than to adult contacts²⁷. Many primary articles and meta-analyses point out the confounding effect of SARS-CoV-2 infections being mostly asymptomatic in young children has on the identification of children as index persons. To some extent, this problem could be avoided in our study since all participants of the event were invited to take part, regardless of age. As all individuals were exposed at the same event and time our study is a very suitable model for the previously described notion, that children are less likely to become infected. Indeed, a recently published meta-analysis by Viner et al. showed a low susceptibility for children and adolescents (OR of 0.56 (95% CI, 0.37-0.85)) which strongly supports our findings of a lower risk of infection in that age group, which is even lower in our study²⁸. Our finding supports the previously shown minor influence on the spreading of the virus by children. The finding that for every 10 additional years of age the risk of infection increases during an event indicates that younger people and their limited role should be considered when measures to contain the pandemic are implemented. It should be mentioned that although children had similar exposure compared to adults and probably spent even less time outside the venue hall, the behaviors of children may be different compared to adults. Therefore, we cannot exclude that our findings of lower seroprevalence in children might be biased by factors very specific to this particular event. Taken together, we demonstrate important risk factors for infection during a superspreading event, which helps to understand transmission dynamics in order to improve comprehensive public health preparedness measures, including mandatory ventilation during indoor events and age-adjusted measures according to different risk of infection.

As to the strengths and limitations of this study, the participant group is extremely well-defined and there was no bias or preselection during enrollment as there was only one

criteria for invitation, namely presence at the event. Because of the time between the event and the study it is possible that participants were infected unrelated to the event, but the official shut down of the community limits this risk. The number of index cases during the event is not known and it is possible that a high number of individuals were already infectious. In addition, the identification of a past SARS-CoV-2 infection via serological test is not perfect and according to the manufacturer their IgG detection is 94.4 % sensitive (on samples collected >10 days after beginning of symptoms or direct detection of virus) and 99.0 % specific (for a ratio ≥ 0.8). For our infection rate analysis this predicts 2 false positives and 10 false negative IgG results. However, when field-tested by the UK National Health Service (NHS) the same assay showed 74.7 % sensitivity (62 false-negatives in our data set) and the same specificity of 99.0 %.

Acknowledgments

We would like to thank the inhabitants of the town of Gangelt for their participation. We would also like to thank the local government of the town where the outbreak took place for their support to conduct the study. Furthermore, we would like to thank the following people who helped with the study: Rudi Kirschen, Gero Wilbring, Janett Wieseler, Marek Korencak, Ryan Natrass, Jernej Pusnik, Maximilian Becker, Ann-Sophie Boucher, Marc Alexander de Boer, Rebekka Dix, Sara Dohmen, Kim Friele, Benedikt Gansen, Jannik Geier, Marie Gronemeyer, Sarah Hundertmark, Nora Jansen, Michael Jost, Louisa Khorsandian, Simon Krzycki, Ekaterina Kuskova, Judith Langen, Silvia Letmathe, Ann-Kathrin Lippe, Jonathan Meinke, Freya Merker, Annika Modemann, Janine Petras, Sophie Marie Porath, Anna Quast, Laurine Reese, Isabel Maria Rehbach, Jonas Richter, Thea Rödig, Eva Schmitz, Tobias Schremmer, Louisa Sommer, Jennifer Speda, Yuhe Tang, Oliver Thanscheidt, Franz Thiele, Johanna Thiele, Julia Tholen, Moritz Transier, Maike van der Hoek, Tillmann Verbeek, Sophia Verspohl, Kira Vordermark, Julian Wirtz, Marina Wirtz, Lisa Zimmer, Philip Koenemann, Adi Yaser, Lisa Anna, Katharina Bartenschlager, Lisa Baum, Roxana Böhmer Romero, Diana de Braganca, Isabelle Engels, Moritz Färber, Carina Fernandez Gonzalez, Lucia Maria Goßner, Victoria Handschuch, Franziska Georgia Liermann, Steffan Meißner, Laura Racenski, Patrick, Denis Raguse, Larissa Reiß, Maximilian Rölle, Franziska Scheele, Chiara Schwippert, Arlene Christin Schwippert, Antonia Seifert, Joshua David Stockhausen, Sofia Waldorf, Leonie Weinhold, Nicolai Trimpop, Julia Reinhardt, Vera Gast, Michelle Yong, Eva Engels, Jonathan Meinke,

Susanne Schmidt, Janine Schulte, Saskia Schmitz, Kübra Bayrak, Regina Frizler, Katarzyna Andryka, Sofía Soler, Bastian Putschli, Thomas Zillinger, Marcel Renn, Patrick Müller, Dillon Corvino, Zeinab Abdullah, Katrin Paeschke, Hiroki Kato, Florian Schmidt, Maximilian Baum, Celina Schüter, Daniel Hinze, Martina Schmidt, Arcangelo Ricchiuto, Sonja Gross, Uta Wolber, Tobias Höller, Marion Zerlett, Esther Sib, Benjamin Marx, Souhaib Aldabbagh. We thank the Medizinischer Dienst der Krankenkasse (MDK) Nordrhein for their generous help in conducting this study. In particular we would like to thank: Tanja Bell, Ina de Hesselle-Taddey, Susanne Goll, Katja Hoffmann-Pruss, Niko Kalamakidis, Verena Kayser, Hildegard Kessler, Norbert Körfer, Daniela Kroll, Florian Messerschmidt, Svenja Peters, Sebastian Schröder, Linda Schroers, Michael Zimmermann, Klaus-Peter Thiele. We thank Stefan Holdenrieder, Alexander Semaan, Bernd Pöttsch, and Georg Nickenig for providing control samples. The idea, the plan, the concept, protocol, the conduct, the data analysis, and the writing of the manuscript of this study were independent of any third parties, including the government of North Rhine-Westphalia, Germany.

Contributors

L.W., E.R., B.S. and H.S. wrote the manuscript. L.W., R.M.S., and E.R. organized and ran the testing center. M.E. inspected the event venue and examined ventilation and air filtration systems. ER and BS organized and performed sample processing, experiments, analyses, and corrected the manuscript. N.L. and A.H. performed statistical analysis. M.C., C.F., and A.K. monitored the study. M.E., K.H.J. and H.S. oversaw the study and corrected the manuscript.

Data availability statement

All data from the study will be made available upon reasonable request.

Funding

The government of North Rhine-Westphalia (Germany) supported the study with 65,000 Euro.

Competing interests

The authors declare no conflict of interest.

References

1. Chakraborty I, Maity P. COVID-19 outbreak: Migration, effects on society, global environment and prevention. *Sci Total Environ* 2020; **728**: 138882.

2. Lloyd-Smith JO, Schreiber SJ, Kopp PE, Getz WM. Superspreading and the effect of individual variation on disease emergence. *Nature* 2005; **438**(7066): 355-9.

3. De Serres G, Markowski F, Toth E, et al. Largest measles epidemic in North America in a decade--Quebec, Canada, 2011: contribution of susceptibility, serendipity, and superspreading events. *J Infect Dis* 2013; **207**(6): 990-8.

4. Shen Z, Ning F, Zhou W, et al. Superspreading SARS events, Beijing, 2003. *Emerg Infect Dis* 2004; **10**(2): 256-60.

5. Streeck H, Schulte B, Kümmerer BM, et al. Infection fatality rate of SARS-CoV2 in a super-spreading event in Germany. *Nat Commun* 2020; **11**(1): 5829.

6. Leclerc QJ, Fuller NM, Knight LE, Funk S, Knight GM. What settings have been linked to SARS-CoV-2 transmission clusters? *Wellcome Open Res* 2020; **5**: 83.

7. Corman VM, Landt O, Kaiser M, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* 2020; **25**(3).

8. Kyriakopoulos AM, Papaefthymiou A, Georgilas N, Doulberis M, Kountouras J. The Potential Role of Super Spread Events in SARS-COV-2 Pandemic; a Narrative Review. *Arch Acad Emerg Med* 2020; **8**(1): e74.

9. Patel KP, Vunnam SR, Patel PA, et al. Transmission of SARS-CoV-2: an update of current literature. *Eur J Clin Microbiol Infect Dis* 2020; **39**(11): 2005-11.

10. Lednicky JA, Lauzardo M, Fan ZH, et al. Viable SARS-CoV-2 in the air of a hospital room with COVID-19 patients. *Int J Infect Dis* 2020; **100**: 476-82.

11. Santarpia JL, Rivera DN, Herrera VL, et al. Aerosol and surface contamination of SARS-CoV-2 observed in quarantine and isolation care. *Sci Rep* 2020; **10**(1): 12732.

12. Günther T, Czech-Sioli M, Indenbirken D, et al. SARS-CoV-2 outbreak investigation in a German meat processing plant. *EMBO Mol Med* 2020; **12**(12): e13296.

13. Pokora R, Kutschbach S, Weigl M, et al. Investigation of superspreading COVID-19 outbreak events in meat and poultry processing plants in Germany: A cross-sectional study. *PLoS One* 2021; **16**(6): e0242456.

14. Nazarenko Y. Air filtration and SARS-CoV-2. *Epidemiol Health* 2020; **42**: e2020049.

15. Vardavas CI, Nikitara K. COVID-19 and smoking: A systematic review of the evidence. *Tob Induc Dis* 2020; **18**: 20.

16. Engin AB, Engin ED, Engin A. Two important controversial risk factors in SARS-CoV-2 infection: Obesity and smoking. *Environ Toxicol Pharmacol* 2020; **78**: 103411.

17. Israel A, Feldhamer E, Lahad A, Levin-Zamir D, Lavie G. Smoking and the risk of COVID-19 in a large observational population study. *medRxiv* 2020: 2020.06.01.20118877.

18. Grundy EJ, Suddek T, Filippidis FT, Majeed A, Coronini-Cronberg S. Smoking, SARS-CoV-2 and COVID-19: A review of reviews considering implications for public health policy and practice. *Tob Induc Dis* 2020; **18**: 58.

19. Changeux JP, Amoura Z, Rey FA, Miyara M. A nicotinic hypothesis for Covid-19 with preventive and therapeutic implications. *C R Biol* 2020; **343**(1): 33-9.

20. Stefano ML, Kream RM, Stefano GB. A Novel Vaccine Employing Non-Replicating Rabies Virus Expressing Chimeric SARS-CoV-2 Spike Protein Domains: Functional Inhibition of Viral/Nicotinic Acetylcholine Receptor Complexes. *Med Sci Monit* 2020; **26**: e926016.

21. Dormoy V, Perotin JM, Gosset P, Maskos U, Polette M, Deslée G. Nicotinic receptors as SARS-CoV-2 spike co-receptors? *Med Hypotheses* 2022; **158**: 110741.
22. F. Oliveira AS, Ibarra AA, Bermudez I, et al. Simulations support the interaction of the SARS-CoV-2 spike protein with nicotinic acetylcholine receptors. *bioRxiv* 2020: 2020.07.16.206680.
23. clinicaltrials.gov. Impact of Smoking and Nicotine on the Risk of Being Infected With COVID-19 (MAGIC). 2021. <https://clinicaltrials.gov/ct2/show/NCT04429815>.
24. Paleiron N, Mayet A, Marbac V, et al. Impact of Tobacco Smoking on the Risk of COVID-19: A Large Scale Retrospective Cohort Study. *Nicotine Tob Res* 2021; **23**(8): 1398-404.
25. Li X, Xu W, Dozier M, He Y, Kirolos A, Theodoratou E. The role of children in transmission of SARS-CoV-2: A rapid review. *J Glob Health* 2020; **10**(1): 011101.
26. Li X, Xu W, Dozier M, et al. The role of children in the transmission of SARS-CoV2: updated rapid review. *J Glob Health* 2020; **10**(2): 021101.
27. Madewell, Z. J., Yang, Y., Longini, I. M., Jr., Halloran, M. E., Dean, N. E. Household Transmission of SARS-CoV-2: A Systematic Review and Meta-analysis. *JAMA Netw Open* 2020; **3**(12): e2031756
28. Viner RM, Mytton OT, Bonell C, et al. Susceptibility to SARS-CoV-2 Infection Among Children and Adolescents Compared With Adults: A Systematic Review and Meta-analysis. *JAMA Pediatr* 2021; **175**(2): 143-56.

Tables

	Not infected N%	Infected N%
Total number	218	186
Female	114 (52%)	89 (48%)
Age:		
<18 years	31 (14%)	15 (8%)
18- 24 years	30 (14%)	20 (11%)
25-39 years	81 (37%)	43 (23%)
40-64 years	71 (33%)	100 (54%)
65+ years	5 (2%)	8 (4%)
BMI [kg/m ²] (Std Dev)	24.3 (5.12)	26.2 (5.16)
Participating household member (Std Dev)	2.1 (1.12)	2.4 (1.16)
Highest level of formal education:		
None	27 (13%)	13 (7%)
lower secondary school	27 (13%)	23 (13%)
secondary school	55 (26%)	71 (39%)
higher education entrance qualification	54 (25%)	34 (18%)
(technical) university degree	52 (24%)	43 (23%)
Duration of attendance [h] (Std Dev)	4.7 (2.06)	5.8 (1.85)
Service team	4 (2%)	22 (12%)
On stage during event	80 (37%)	62 (34%)
Member of „Council of 11“	6 (3%)	18 (10%)
On stage during “Finale”	26 (12%)	48 (26%)
Behavior during break:		
Remaining seated	73 (36%)	85 (48%)

	Going outside	114 (55%)	72 (39%)
Alcohol consumption [drink] (Std Dev)		11.3 (7.76)	12.2 (7.40)
Former smoker		34 (16%)	45 (24%)
Active smoker (≥10 cigarettes/day)		54 (25%)	23 (12%)
At least one comorbidity		29 (13%)	28 (15%)
Avg. distance to other participants [m] (Std Dev)		9.2 (1.68)	9.1 (1.70)
Distance to air inlet [m] (Std Dev)		6.1 (3.22)	6.0 (3.30)
Distance to air outlet [m] (Std Dev)		4.8 (2.94)	5.1 (2.87)

Table 1: Distribution of demographic factors and exposure information of interest among study participants who tested positive or negative in serology test of SARS-Cov2-infection

‘Council of 11’ stands for the hosts of the events located on stage (personnel switched during the break).
‘Finale’ describes the final presentation of the event with all performers on stage.

Figure Legends

Figure 1: Study participants

Enrollment and flow of participants through the study. Downstream sample processing included centrifugation of blood samples for plasma collection (SARS-CoV-2 ELISAs), and viral RNA extraction from swab samples (SARS-CoV-2 RT-PCR).

Figure 2: Odds ratios for the likelihood of SARS-CoV-2 infection by age groups

Participants were divided into age groups of 8, 15, or 25 years, participants younger than 18 or older than 65 years. Participants were considered to have been infected during the event if they were SARS-CoV-2 antibody positive (ELISA).

Figure 3: Reconstructed 3D-Model of the venue hall

The venue was a single open space with a stage on one end and a bar as well as the exit on the opposite end. Distribution of tables and seating was as indicated by table and chairs symbols. Please note that the people pictured are illustrative and do not represent individual participants. Self-administered questionnaires included questions about main seating-position of the participants during the evening event as specifying table and seat with the help of a schematic seating plan. Metric room coordinates for all tables, seats, and ventilation-points were assessed and the seating was reconstructed from pictures taken during the event. Therefore, the location of the stage, the bar, the exit as well as the tables and the air-inlets/outlets were reconstructed in a 3D-Model. The original external dimensions of the building were 27m x 13.20m x 4.20m. Tables, where more than 7 infected individuals have stayed are colored in dark red, this includes the stage and bar as well. Air-inlets are colored in violet and the air-outlets in blue. Infected participants had been seated mostly at tables close to the bar, the bar itself and on stage. One table with 8 out of 11 infected people, was located far away from the bar at the other side of the hall and close to an air inlet. The group sitting on stage showed as well high numbers of infection (18 infected out of 24).

Figure 4: Odds ratios for the association of SARS-CoV-2 infection with specific activities of the participants and their location in the venue relative to ventilation shafts

The model was additionally adjusted for age, sex, duration of attendance, participation in multiple activities, and cumulative proximity to other infected persons, and common household.

Figure 5: Odds ratios for symptoms of SARS-CoV-2 antibody-positive participants in the 14 days following the super spreading event

The information on symptoms was derived from the self-administered questionnaire, which was filled out on the day of sample collection. Odds ratio estimates (OR) are shown with confidence intervals.




Fig. 1: Study participants. Enrollment and flow of participants through the study. Downstream sample processing included centrifugation of blood samples for plasma collection (SARS-CoV-2 ELISAs), and viral RNA extraction from swab samples (SARS-CoV-2 RT-PCR).

338x123mm (78 x 78 DPI)




Fig. 2 Odds-Ratio for the likelihood of SARS-CoV-2 infection by age groups. Participants were divided into age groups of 8, 15, or 25 years, participants younger than 18 or older than 65 years. Participants were considered to have been infected during the event if they were SARS-CoV-2 antibody positive (ELISA).

283x160mm (300 x 300 DPI)




Fig. 3: Reconstructed 3D-Model of the venue hall. The venue was a single open space with a stage on one end and a bar as well as the exit on the opposite end. Distribution of tables and seating was as indicated by table and chairs symbols. Please note that the people pictured are illustrative and do not represent individual participants. Self-administered questionnaires included questions about main seating-position of the participants during the evening event as specifying table and seat with the help of a schematic seating plan. Metric room coordinates for all tables, seats, and ventilation-points were assessed and the seating was reconstructed from pictures taken during the event. Therefore, the location of the stage, the bar, the exit as well as the tables and the air-inlets/outlets were reconstructed in a 3D-Model. The original external dimensions of the building were 27m x 13.20m x 4.20m. Tables, where more than 7 infected individuals have stayed are colored in dark red, this includes the stage and bar as well. Air-inlets are colored in violet and the air-outlets in blue. Infected participants had been seated mostly at tables close to the bar, the bar itself and on stage. One table with 8 out of 11 infected people, was located far away from the bar at the other side of the hall and close to an air inlet. The group sitting on stage showed as well high numbers of infection (18 infected out of 24).

155x96mm (300 x 300 DPI)




Fig. 4: Odds ratios for the association of SARS-CoV-2 infection with specific activities of the participants and their location in the venue relative to ventilation shafts. The model was additionally adjusted for age, sex, duration of attendance, participation in multiple activities, and cumulative proximity to other infected persons, and common household.

259x115mm (120 x 120 DPI)






Fig. 5: Odds ratios for symptoms of SARS-CoV-2 antibody-positive participants in the 14 days following the super spreading event. The information on symptoms was derived from the self-administered questionnaire, which was filled out on the day of sample collection. Odds ratio estimates (OR) are shown with confidence intervals.

280x208mm (120 x 120 DPI)


Supplementary Material



Suppl. Fig. 1: Correlation of SARS-CoV-2 Euroimmun ELISA results for IgA and IgG. The correlation of IgA levels to IgG levels in the same person was significant (r: Pearson coefficient, $p<0.0001$, 95 % CI, 0.7043 to 0.7902). The dotted lines mark the ratios above which each ELISA result is considered positive.



Suppl. Fig. 2: Correlation of plasma neutralization capacity and IgG ELISA results (Euroimmun) from each donor. The dotted line marks the ratio above which the ELISA result is considered positive. The correlation coefficient (Pearson) was 0.3667 (95 % CI, 0.2275 to 0.4192, $p<0.0001$). Samples with a negative result in the neutralization assay were set as 0.1 here so as to appear on the logarithmic axis.



Supplemental Figure 3: Quantile plot of observed p-values from analyses of inverse distance [1/m] to single specific study participants as risk factor for corona-virus infection. In case of no association, the ordered log-transformed p-values are expected to lie on, or below the diagonal. Panel A: results from crude analyses, Panel B: analyses were adjusted for age, sex, common household and duration of attendance.

	OR	95% confidence interval		p-value
proximity to infected persons [sum 1/m]	0,99	0,98	1,01	0,430
adjusted a)	1,00	0,98	1,02	0,957
mutually adjusted b)	0,99	0,97	1,01	0,571
mutually adjusted c)	0,99	0,97	1,02	0,646
Alternative consideration in distance-bands				
Infected persons within ≤1.5m [count]	1,01	0,96	1,07	0,681
Infected persons in 1.5 - ≤3m [count]	0,96	0,92	1,00	0,043
Infected persons in 3 - ≤4.5m [count]	1,03	1,00	1,06	0,023
Infected within ≤1.5m [count] adjusted a)	1,03	0,97	1,10	0,366
Infected in 1.5 - ≤3m [count]	0,96	0,92	1,01	0,113
Infected in 3 - ≤4.5m [count]	1,03	1,00	1,06	0,083
Infected within ≤1.5m [count] mutually adjusted b)	1,01	0,95	1,07	0,734
Infected in 1.5 - ≤3m [count]	0,98	0,94	1,02	0,359
Infected in 3 - ≤4.5m [count]	1,05	1,02	1,08	0,001
Infected within ≤1.5m [count] mutually adjusted c)	1,02	0,95	1,09	0,638
Infected in 1.5 - ≤3m [count]	0,98	0,93	1,03	0,363
Infected in 3 - ≤4.5m [count]	1,04	1,00	1,07	0,041

Supplementary table 1: Estimated relative risk of SARS-CoV-2 infection (IGG-positive) from logistic regression on summary measures of spatial proximity between participants in terms of odds ratio estimates (OR) with confidence interval and p-values. a) adjusted for sex, age, common household and duration. b) multivariate analysis, mutually adjusted for distance to ventilation system, participation in (multiple) performances, going out of doors during the intermission, and participating in the grand finale. c) multivariate analysis, mutually adjusted for distance to ventilation system, participation in (multiple) performances, going out of doors during the intermission, and participating in the grand finale and adjusted for sex, age, common household and duration.

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	3
Objectives	3	State specific objectives, including any prespecified hypotheses	3
Methods			
Study design	4	Present key elements of study design early in the paper	3,4
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	3-7
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants	3,4
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	2
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	4
Bias	9	Describe any efforts to address potential sources of bias	3
Study size	10	Explain how the study size was arrived at	3
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	4
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	4
		(b) Describe any methods used to examine subgroups and interactions	5
		(c) Explain how missing data were addressed	4
		(d) If applicable, describe analytical methods taking account of sampling strategy	
		(e) Describe any sensitivity analyses	14
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	19
		(b) Give reasons for non-participation at each stage	19
		(c) Consider use of a flow diagram	19
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	7
		(b) Indicate number of participants with missing data for each variable of interest	7
Outcome data	15*	Report numbers of outcome events or summary measures	7,8
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	9,10

		(b) Report category boundaries when continuous variables were categorized	8,10
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	9-11
Discussion			
Key results	18	Summarise key results with reference to study objectives	11-13
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	14
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	13,14
Generalisability	21	Discuss the generalisability (external validity) of the study results	11
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	16

*Give information separately for exposed and unexposed groups.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.